

On a Unified Theory of Generalized Branes Coupled to Gauge Fields, Including the Gravitational and Kalb-Ramond Fields

M. Pavšič

Jožef Stefan Institute, Jamova 39, SI-1000 Ljubljana, Slovenia; E-mail:

matej.pavsic@ijs.si

ABSTRACT

We investigate a theory in which fundamental objects are branes described in terms of higher grade coordinates $X^{\mu_1 \dots \mu_n}$ encoding both the motion of a brane as a whole, and its volume evolution. We thus formulate a dynamics which generalizes the dynamics of the usual branes. Geometrically, coordinates $X^{\mu_1 \dots \mu_n}$ and associated coordinate frame fields $\{\gamma_{\mu_1 \dots \mu_n}\}$ extend the notion of geometry from spacetime to that of an enlarged space, called Clifford space or C -space. If we start from 4-dimensional spacetime, then the dimension of C -space is 16. The fact that C -space has more than four dimensions suggests that it could serve as a realization of Kaluza-Klein idea. The “extra dimensions” are not just the ordinary extra dimensions, they are related to the volume degrees of freedom, therefore they are physical, and need not be compactified. Gauge fields are due to the metric of Clifford space. It turns out that amongst the latter gauge fields there also exist higher grade, antisymmetric fields of the Kalb-Ramond type, and their non-Abelian generalization. All those fields are naturally coupled to the generalized branes, whose dynamics is given by a generalized Howe-Tucker action in curved C -space.

1 Introduction

Point particle is an idealization never found in nature. Physical objects are extended and possess in principle infinitely many degrees of freedom. It is now widely accepted that even at the “fundamental” level objects are extended. Relativistic strings and higher dimensional extended objects, branes, have attracted much attention during last three decades [1, 2, 3, 4].

An extended object, such as a brane, during its motion sweeps a worldsheet, whose points form an n -dimensional manifold¹ V_n embedded in a target space(time) V_N . Worldsheet is usually considered as being formed by a set of *points*, that is, with a worldsheet we associate a *manifold of points*, V_n . Alternatively, we can consider a worldsheet as being formed by a set of closed $(n - 1)$ -branes (that we shall call “loops”). For instance, a string world sheet V_2 can be considered as being formed by a set of 1-loops. In particular such a 1-loop can be just a closed string which in the course of its evolution sweeps a worldsheet V_2 which, in this case, has the form of a world tube. But in general, this need not be the case. A set of 1-loops on V_2 need not coincide with a family of strings for various values a time-like parameter. Thus even a worldsheet swept by an open string can be considered as a set of closed loops. The ideas that we pursue here are motivated and based to certain extent by those developed in refs. [6]–[8]. We shall employ the very powerfull geometric language based on Clifford algebra [9, 10], which has turned out to be very suitable for an elegant formulation of p -brane theory and its generalization [11]–[16]. We will employ the property that multivectors of various definite grades, i.e., R -vectors, since they represent oriented lines, areas, volumes,..., shortly, R -volumes (that we will also call R -areas), can be used in the description of branes. With a brane one can associate an oriented R -volume (R -area). Superpositions of R -vectors are generic Clifford numbers, that we call polyvectors. They represent geometric objects, which are superposition of oriented lines, areas, volumes,..., that we associate, respectively, with point particles, closed strings, closed 2-branes,... , or alternatively, with open strings, open 2-branes, open 3-branes, etc.. A polyvector is thus used for description of a physical object, a *generalized brane*, whose components are branes of various dimensionalities.

We thus describe branes by means of higher grade coordinates $x^{\mu_1 \dots \mu_R}$, $R = 0, 1, 2, \dots, N$, corresponding to an oriented R -area associated with a brane, where N is the dimension of the spacetime V_N we started from. The latter coordinates are *collective coordinates*, [13, 14], analogous to the center of mass coordinates [11]. They do not provide a full description of an extended object, they merely sample it. Nevertheless, if

¹In the literature, the name ‘worldsheet’ is often reserved for a 2-dimensional surface swept by 1-dimensional string. Here we use ‘worldsheet’ for a surface of any dimension n swept by an $(n - 1)$ -dimensional brane. By symbols V_n and V_N we denote *manifolds* (we adopt an old practice), and *not* vectors spaces.

higher grade coordinates $x^{\mu_1 \dots \mu_R}$ are given, then certainly we have more information about an extended object than in the case when only its center of mass coordinates are given. By higher grade coordinates we no longer approximate an extended object with a point like object; we take into account its extra structure.

We associate all those higher grade coordinates with *points* of an 2^N -dimensional space, called *Clifford space*, shortly *C-space*, denoted C_{V_N} . Every point of C_{V_N} represents a possible *extended event*, associated with a generalized brane.

In order to consider an object's dynamics, one has to introduce a continuous parameter, say τ , and consider a mapping $\tau \rightarrow x^{\mu_1 \dots \mu_R} = X^{\mu_1 \dots \mu_R}(\tau)$. So functions $X^{\mu_1 \dots \mu_R}(\tau)$ describe a curve in an 2^N -dimensional space C_{V_N} . This generalizes the concept of *worldline* $X^\mu(\tau)$ in spacetime V_N . The action principle is given by the *minimal length action* in C_{V_N} . That the objects, sampled by $X^{\mu_1 \dots \mu_R}$ satisfy such dynamics is our *postulate* [11, 12, 16, 15], we do *not* derive it.

The intersection of a *C-space* worldline $X^{\mu_1 \dots \mu_R}(\tau)$ with an underlying spacetime V_N (which is a subspace of C_{V_N}) gives, in general an extended event². Therefore, what we observe in spacetime are “instantonic” extended objects that are localized both in space-like and time-like directions³. According to this generalized dynamics, worldlines are infinitely extended in C_{V_N} , but in general, their intersections with subspace V_N are finite. In spacetime V_N we observe finite objects whose time like extension may increase with evolution, and so after a while they mimic the worldlines of the usual relativity theory. This has been investigated in refs. [11, 15]. We have also found that such *C-space* theory includes the Stueckelberg theory [17]–[24] as a particular case, and also has implications for the resolution of the long standing problem of time in quantum gravity [25, 12, 26].

Objects described by coordinates $x^{\mu_1 \dots \mu_R}$ are *points* in Clifford space C_{V_N} , also called *extended events*. Objects given by functions $X^{\mu_1 \dots \mu_R}(\tau)$ are *worldlines* in C_{V_N} . A further possibility is to consider, e.g., continuous sets of extended events, described by functions $X^{\mu_1 \dots \mu_R}(\xi^A)$, $A = 1, 2, \dots, 2^n$, $n < N$, where $\xi^A \equiv \xi^{a_1 \dots a_r}$, $r = 0, 1, 2, \dots, n < N$, are 2^n higher grade coordinates denoting oriented r -areas in the parameter space $\mathbb{R}^n$. Functions $X^{\mu_1 \dots \mu_R}(\xi^A)$ describe a 2^n -dimensional surface in C_{V_N} . This generalizes the concept of *worldsheet* or *world manifold* $X^\mu(\xi^a)$, $a = 1, 2, \dots, n$, i.e., the surface that an evolving brane sweeps in the embedding spacetime V_N .

A *C-space* worldline $X^{\mu_1 \dots \mu_R}(\tau)$ does not provide a “full” description of an extended object, because not “all” degrees of freedom are taken into account; $X^{\mu_1 \dots \mu_R}(\tau)$ only provides certain “collective” degrees of freedom that sample an extended object. On the contrary, a *C-space* worldsheet $X^{\mu_1 \dots \mu_R}(\xi^A)$ provides much more detailed description, be-

²Analogously, in spacetime the intersection of an ordinary worldline with a 3-dimensional slice gives a point.

³ They are the analog of $p = -1$ branes (instantons) that are important in string theories.

cause of the presence of 2^n continuous parameters ξ^A , on which the generalized coordinate functions $X^{\mu_1 \dots \mu_R}(\xi^A)$ depend. In particular, the latter functions can be such that they describe just an ordinary worldsheet, swept by an ordinary brane. But in general, they describe more complicated extended objects, with an extra structure.

We equip our manifold C_{V_N} with metric, connection and curvature. In the case of *vanishing curvature*, we can proceed as follows. We choose in C_{V_N} an *origin* $\mathcal{E}_0$ with coordinates $X^{\mu_1 \dots \mu_R}(\mathcal{E}_0) = 0$. This enables us to describe points $\mathcal{E}$ of C_{V_N} with *vectors* pointing from $\mathcal{E}_0$ to $\mathcal{E}$. Since those vectors are Clifford numbers, we call them *polyvectors*. So points of our *flat* space C_{V_N} (i.e., with vanishing curvature) are described by polyvectors $x^{\mu_1 \dots \mu_R} \gamma_{\mu_1 \dots \mu_R}$, where $\gamma_{\mu_1 \dots \mu_R}$ are basis Clifford numbers, that span a Clifford algebra $\mathcal{C}_N$.

So our extended objects, the events $\mathcal{E}$ in C_{V_N} , are described by Clifford numbers. This actually brings *spinors* into the description, since, as is well known, the elements of left (right) ideals of a Clifford algebra represent spinors [27]. So one does not need to postulate spinorial variables separately, as is usually done in string and brane theories. Our model is an alternative to the theory of spinning branes and supersymmetric branes, including spinning strings and superstrings [1]. In refs. [16] it was shown that the 16-dimensional Clifford space provides a framework for a consistent string theory. One does not need to postulate extra dimensions of spacetime. One can start from 4-dimensional spacetime, and finds that the corresponding Clifford space provides enough degrees of freedom for a string, so that the Virasoro algebra has no central charges. According to this theory all 16 dimensions of Clifford space are physical and thus observable [11, 28, 29], because they are related to the extended nature of objects. Therefore, there is no need for compactification of the extra dimensions of Clifford space.

As a next step it was proposed [28, 29] that curved 16-dimensional Clifford space can provide a realization of Kaluza-Klein theory. Gravitational as well as other gauge interactions can be unified within such a framework. In ref. [28, 29] we considered Yang-Mills gauge field potentials as components of the C -space connection, and Yang-Mills gauge field strengths as components of the curvature of that connection. It was also shown [29] that in a curved C -space which admits K isometries, Yang-Mills gauge potentials occur not only in the connection, but also in the metric, or equivalently, in the vielbein. In this paper we concentrate on the latter property, and further investigate it by studying the brane action in curved background C -space. So we obtain the minimal coupling terms in the classical generalized brane action, and we show that the latter coupling terms contain the ordinary 4-dimensional gravitational fields, Yang-Mills gauge fields A_μ^α , $\alpha = 1, 2, \dots, K$, and also the higher grade, in general non-Abelian, gauge fields $A_{\mu_1 \dots \mu_R}^\alpha$ of the Kalb-Ramond type. We thus formulate an elegant, unified theory for the classical generalized branes coupled to all those various fields.

2 On the description of extended objects

2.1 Worldsheet described by a set of point events

As an example of a relativistic extended object, let us first consider the string. An evolving string sweeps a worldsheet, a physical object in the embedding spacetime. Worldsheet can be considered as being formed by a set of point events. So with a worldsheet we can associate a 2-dimensional manifold V_2 , called *world manifold*, embedded in an N -dimensional target space V_N . With every *point* on V_2 we associate two parameters (coordinates) ξ^a , $a = 0, 1$ which are arbitrary (like “house numbers”). The embedding of V_2 into V_N is described by the mapping

$$\xi^a \rightarrow x^\mu = X^\mu(\xi^a), \quad \xi^a \in R^n \subset \mathbb{R}^n \quad (1)$$

where x^μ , $\mu = 0, 1, 2, \dots, N-1$ are coordinates describing position in V_N , whilst $X^\mu(\xi^a)$ are embedding functions (Fig.1), defined over a domain R^n within a set $\mathbb{R}^n$ of real numbers.




Figure 1: A worldsheet can be considered as being formed by a set of point events associated with *points* of a world manifold V_2 embedded in V_N , described by the mapping $\xi^a \rightarrow x^\mu = X^\mu(\xi^a)$.

With our worldsheet we thus associate a 2-parameter set of *points* described by functions $X^\mu(\xi^0, \xi^1)$. We distinguish here the physical object, the worldsheet, from the corresponding mathematical object, the manifold (world manifold) V_2 . Strictly, we should use two different symbols for those different objects. In practice, we will not be so rigorous, and we will simply denote worldsheet by the symbol V_2 (in general V_N).

2.2 Worldsheet described by a set of loops

In previous section a worldsheet was described by a 2-parameter set of points described by functions $X^\mu(\xi^0, \xi^1)$. Alternatively, instead of points we can consider closed lines, loops, each being described by functions $X^\mu(s)$, where $s \in [s_1, s_2] \subset \mathbb{R}$ is a parameter along a loop. A 1-parameter family of such loops $X^\mu(s, \alpha)$, $\alpha \in [\alpha_1, \alpha_2] \subset \mathbb{R}$ sweeps a worldsheet V_2 . This holds regardless of whether such worldsheet is open or closed. However, in the case of an open worldsheet, the loops are just kinematically possible objects, and they cannot be associated with physical closed strings. In the case of a closed worldsheet, a world tube, we can consider it as being swept by an evolving closed string.

We will now demonstrate, how with every loop one can associate an oriented area, whose projections onto the coordinate planes are $X^{\mu\nu}$. The latter quantities are functionals of a loop $X^\mu(s)$. If we consider not a single loop, but a family of loops $X^\mu(s, \alpha)$, $\alpha \in [\alpha_1, \alpha_2]$, then $X^{\mu\nu}$ are *functions of parameter* α , besides being *functionals of a loop*. So we obtain a 1-parameter family of oriented areas described by functions $X^{\mu\nu}(\alpha)$. Let us stress again that for every *fixed* α , it holds, of course, that $X^{\mu\nu}$ are functionals of $X^\mu(s, \alpha)$.

If we choose a loop B on V_2 , i.e., a loop from a given family $\{X^\mu(s, \alpha), \alpha \in [\alpha_1, \alpha_2]\}$, then we obtain the corresponding components $X^{\mu\nu}$ of the oriented area by performing the integration of infinitesimal oriented area elements over a chosen surface whose boundary is our loop B . Given a boundary loop B , it does not matter which surface we choose. In the following, for simplicity, we will choose just our worldsheet V_2 for the surface.

Let us now consider a surface element on V_2 . Let $d\xi_1 = d\xi_1^a e_a$ and $d\xi_2 = d\xi_2^a e_a$ be two infinitesimal vectors on V_2 , expanded in terms of basis vectors e_a , $a = 0, 1$. An infinitesimal oriented area is given by the wedge product

$$d\Sigma = d\xi_1 \wedge d\xi_2 = d\xi_1^a d\xi_2^b e_a \wedge e_b = \frac{1}{2} d\xi^{ab} e_a \wedge e_b \quad (2)$$

where

$$d\xi^{ab} = d\xi_1^a d\xi_2^b - d\xi_2^a d\xi_1^b \quad (3)$$

At every point $\xi \in V_2$ basis vectors e_a span a 2-dimensional linear vector space, a tangent space $T_\xi(V_2)$. Following an old tradition (see, e.g., [30, 31]) we use symbol V_n for an n -dimensional surface embedded in an N -dimensional space V_N . Thus V_n , and in particular V_2 , denotes a manifold, and not a vector space. In order to simplify our wording, an expression like “vectors e_a on V_2 ” will mean “tangent vectors e_a at a point $\xi \in V_2$ ”. So whenever we talk about vectors, or whatever geometric objects, on a manifold (or in a manifold) we just mean that to a given point of the manifold we attach a geometric object (see, e.g., [32]). The latter object, of course, is not an element of our manifold, but of the tangent space. Basis vectors on V_2 can be considered as being induced from the target

space basis vectors γ_μ :

$$e_a = \partial_a X^\mu \gamma_\mu \quad (4)$$

In the following we will adopt the geometric calculus in which basis vectors are *Clifford numbers* satisfying

$$\gamma_\mu \cdot \gamma_\nu \equiv \frac{1}{2}(\gamma_\mu \gamma_\nu + \gamma_\nu \gamma_\mu) = g_{\mu\nu} \quad (5)$$

where $g_{\mu\nu}$ is the metric of V_N . Eq. (5) defines the *inner product* of two vectors as the symmetric part of the Clifford product $\gamma_\mu \gamma_\nu$. The antisymmetric part of $\gamma_\mu \gamma_\nu$ is identified with the *wedge* or *outer product*

$$\gamma_\mu \wedge \gamma_\nu \equiv \frac{1}{2}(\gamma_\mu \gamma_\nu - \gamma_\nu \gamma_\mu) \quad (6)$$

Analogous relations we have for the worldsheet basis vectors e_a :

$$e_a \cdot e_b \equiv \frac{1}{2}(e_a e_b + e_b e_a) = \gamma_{ab} \quad (7)$$

$$e_a \wedge e_b \equiv \frac{1}{2}(e_a e_b - e_b e_a) \quad (8)$$

where γ_{ab} is the metric on V_2 which, according to eq.(4), can be considered as being induced from the target space.

If we insert the relation (4) into eq.(2) we have

$$d\Sigma = \frac{1}{2} d\xi^{ab} \partial_a X^\mu \partial_b X^\nu \gamma_\mu \wedge \gamma_\nu \quad (9)$$

This is an infinitesimal *bivector* or *2-vector* in the target space V_N .

A finite 2-vector is obtained upon integration⁴ over a finite surface Σ_B enclosed by a loop B :

$$\begin{aligned} \int_{\Sigma_B} d\Sigma &\equiv \frac{1}{2} X^{\mu\nu} \gamma_\mu \wedge \gamma_\nu = \frac{1}{2} \int_{\Sigma_B} d\xi^{ab} \partial_a X^\mu \partial_b X^\nu \gamma_\mu \wedge \gamma_\nu \\ &= \frac{1}{2} \int_{\Sigma_B} d\xi^{ab} \frac{1}{2} (\partial_a X^\mu \partial_b X^\nu - \partial_a X^\nu \partial_b X^\mu) \gamma_\mu \wedge \gamma_\nu \end{aligned} \quad (10)$$

From eq.(10) we read

$$X^{\mu\nu}[B] = \frac{1}{2} \int_{\Sigma_B} d\xi^{ab} (\partial_a X^\mu \partial_b X^\nu - \partial_a X^\nu \partial_b X^\mu) \quad (11)$$

By Stokes theorem this is equal to

$$X^{\mu\nu}[B] = \frac{1}{2} \oint_B ds \left(X^\mu \frac{\partial X^\nu}{\partial s} - X^\nu \frac{\partial X^\mu}{\partial s} \right) \quad (12)$$

⁴Such integration poses no problem in *flat* V_N . In curved space we may still use the same expression (10) which then *defines* such integral that all vectors are carried together by means of parallel transport along geodesics into a chosen point of V_N where the integration is actually performed [33].

where $X^\mu(s)$ describes a boundary loop B , s being a parameter along the loop.

Eq.(12) demonstrates that $X^{\mu\nu}$ are components of the bivector, determining an oriented area, associated with a surface enclosed by a loop $X^\mu(s)$ on the worldsheet V_2 . Hence there is a close correspondence between surfaces and the boundary loops. The components $X^{\mu\nu}$ can be therefore be considered as bivector coordinates of a loop. These are collective coordinates, since the detailed shape (configuration) of the loop is not determined by $X^{\mu\nu}$. Only the oriented area associated with a surface enclosed by the loop is determined by $X^{\mu\nu}$. Therefore $X^{\mu\nu}$ refers to a class of loops, from which we may choose a *representative loop*, and say that $X^{\mu\nu}$ are its coordinates. From now on, ‘loop’ we will be often a short hand expression for a *representative loop* in the sense above.

By means of eqs. (10)–(12) we have performed a mapping from an *infinite dimensional space* of loops $X^\mu(s)$ into a *finite dimensional space* of oriented areas $X^{\mu\nu}$. Instead of describing loops by infinite dimensional objects $X^\mu(s)$, we can describe them by finite dimensional objects, oriented areas, with bivector coordinates $X^{\mu\nu}$. *We have thus arrived at a finite dimensional description of loops* (in particular, closed strings), the so called *quenched minisuperspace description* suggested by Aurilia et al. [13].

When we consider not a single loop $X^\mu(s)$, but a 1-parameter family of loops $X^\mu(s, \alpha)$, we have a worldsheet, considered as being formed by a set of loops. By means of eqs. (10)–(12), with every loop within such a family, i.e., for a fixed α , we can associate bivector coordinates $X^{\mu\nu}$. For variable α we then obtain functions $X^{\mu\nu}(\alpha)$. This is a quenched minisuperspace description of a worldsheet. A full description is in terms of embedding functions $X^\mu(\xi^0, \xi^1)$, or a family of loops $X^\mu(s, \alpha)$.

In the following we will consider two particular choices for parameter α .

In eq.(10) we have the expression for an oriented area associated with a loop B . It has been obtained upon the integration of the infinitesimal oriented surface elements (2). Besides the oriented area we can associate with our loop on V_2 also a *scalar* quantity, namely the *scalar area* $\mathcal{A}$ which we obtain according to

$$\mathcal{A} = \int_{\Sigma_B} \sqrt{d\Sigma^\dagger \cdot d\Sigma} \quad (13)$$

Here ‘ $\dagger$ ’ denotes *reversion*, that is the operation which reverse the order of vectors in a product. Using the relation $e_a \wedge e_b = e \epsilon_{ab}$, where $e = e_1 \wedge e_2$ is the pseudoscalar in 2-dimensional space V_2 such that $e^\dagger \cdot e = (e_2 \wedge e_1) \cdot (e_1 \wedge e_2) = \gamma_{11}\gamma_{22} - \gamma_{21}\gamma_{12} = \det \gamma_{ab} \equiv \gamma$, we find

$$d\Sigma^\dagger \cdot d\Sigma = \frac{1}{4} \gamma (d\xi^{ab} \epsilon_{ab})^2 \quad (14)$$

i.e.

$$\mathcal{A} = \int \sqrt{d\Sigma^\dagger \cdot d\Sigma} = \frac{1}{2} \int \sqrt{|\gamma|} d\xi^{ab} \epsilon_{ab} = \int \sqrt{|\gamma|} d\xi^{12} \quad (15)$$

If we choose $d\xi_1^a = (d\xi^1, 0)$, $d\xi_2^a = (0, d\xi^2)$, then $d\xi^{12} = d\xi^1 d\xi^2$.

We imagine that our surface V_2 is covered by a family of loops $X^\mu(\alpha, s)$ (Fig.2), such that the totality of points of all those loops is in one-to-one correspondence with the points of the manifold V_2 , and that for parameter α we take the scalar area $\mathcal{A}$. To every loop there belong bivector coordinates $X^{\mu\nu}$ (calculated according to eq.(10)) and a scalar parameter $\mathcal{A}$ (calculated according to eq. (15)), determining the the scalar area. Dependence of $X^{\mu\nu}$ on $\mathcal{A}$ is characteristic for a given class of surfaces V_2 . If we consider a different class of surface V_2 , then functions $X^{\mu\nu}(\mathcal{A})$ are in general different (Fig. 3)




Figure 2: We consider a loop on V_2 . It determines an *oriented area* whose extrinsic 2-vector coordinates are $X^{\mu\nu}$. The *scalar area* of the surface element enclosed by the loop is $\mathcal{A}$. Given an initial loop, functions $X^{\mu\nu}(\mathcal{A})$ are characteristic for a class of the surface V_2 .




Figure 3: (a) Examples of two different surfaces belonging to a class of surfaces that all satisfy equation $X^{12}(\mathcal{A}) = k$. Constant k differs from zero, if a cillindric surface has only one boundary loop, so that, e.g., the upper part is open, whilst the lower part is closed. (b) Example of surfaces belonging to a class of surfaces that satisfy equation $X^{12}(\mathcal{A}) = k\mathcal{A}$.

Instead of starting with a given surface V_2 on which we determine a family of loops and calculate the functions $X^{\mu\nu}(\mathcal{A})$, we can start from the other end. We may assume that all what is known are functions $X^{\mu\nu}(\mathcal{A})$. From those functions we do not know what

the surface (worldsheet) V_2 exactly is, but we have some partial information (see Fig. 3), up to a class of surfaces (worldsheets). Functions $X^{\mu\nu}(\mathcal{A})$ provide a means of describing a surface V_2 , although not in all details.

Instead of the scalar area $\mathcal{A}$ we may take as the parameter any other parameter⁵. We may take, for instance, just the integral⁶ in the parameter space $\mathbb{R}^2$

$$\xi^{ab}(B) = \int_{\Sigma_B} d\xi'^{ab} \quad (16)$$

taken over a domain corresponding to a surface Σ_B with boundary B .

So we have a mapping $B \rightarrow \xi^{ab}(B)$, such that to any boundary loop B of our family there correspond parameters ξ^{ab} . Because of the property $\xi^{ab} = -\xi^{ba}$, (where $a, b = 0, 1$ if V_2 is time like), there is in fact a single parameter ξ^{12} . The extrinsic 2-vector coordinates $X^{\mu\nu}$ are functions of ξ^{ab} . The mapping

$$\xi^{ab} \rightarrow x^{\mu\nu} = X^{\mu\nu}(\xi^{ab}) \quad (17)$$

determines a class of surfaces V_2 , embedded in V_N , which are all in accordance with eq. (17). Knowing the functions $X^{\mu\nu}(\xi^{ab})$ means knowing a class $\{V_2\}$, but not a particular $V_2 \in \{V_2\}$.

The mapping $\tau \rightarrow x^\mu = X^\mu(\tau)$, involving vector coordinates, describes a curve (a “worldline”) in the space spanned by vectors γ_μ . The derivative of X^μ with respect to τ , i.e., $\dot{X}^\mu = dX^\mu/d\tau$, is the tangent vector to the worldline, or *velocity*.

Similarly, the mapping (17), involving bivector coordinates, describes a curve (a “worldline”) in the space spanned by the bivectors $\gamma_\mu \wedge \gamma_\nu$, and we can calculate the derivative

$$\partial_{ab} X^{\mu\nu} \equiv \frac{\partial X^{\mu\nu}}{\partial \xi^{ab}} \quad (18)$$

which generalizes the concept of velocity.

2.3 Generalization to arbitrary dimensions

Let us now consider extended objects associated with manifolds V_n that have arbitrary dimension n and are embedded in a target space V_N of dimension N . An infinitesimal infinitesimal oriented area element on V_n is an n -vector

$$d\Sigma = d\xi_1 \wedge d\xi_2 \wedge \dots \wedge d\xi_n = d\xi_1^{a_1} d\xi_2^{a_2} \dots d\xi_n^{a_n} e_{a_1} \wedge e_{a_2} \wedge \dots \wedge e_{a_n} = \frac{1}{n!} d\xi^{a_1 \dots a_n} e_{a_1} \wedge \dots \wedge e_{a_n} \quad (19)$$

⁵In a suitable choice of parameters ξ^a the determinant of the metric γ_{ab} on V_2 can be constant, that is $\partial_a |\gamma| = 0$. Choosing $|\gamma| = 1$, the scalar area $\mathcal{A}$ is given just by the integral (16).

⁶Similarly, in describing a world line $X^\mu(\tau)$, we may take any parameter τ which, in particular, can be the length of the worldline. The analog of eq. (16) is $\tau = \int_0^\tau d\tau'$.

where

$$d\xi^{a_1 \dots a_n} = d\xi_1^{a_1} d\xi_2^{a_2} \dots d\xi_n^{a_n} \quad (20)$$

If we consider the basis vectors e_a on V_n as being induced from the basis vectors γ_μ of the embedding space V_N according to the relation (4), we have

$$d\Sigma = \frac{1}{n!} dX^{\mu_1 \dots \mu_n} \gamma_{\mu_1} \wedge \dots \wedge \gamma_{\mu_n} = \frac{1}{n!} d\xi^{a_1 \dots a_n} \partial_{a_1} X^{\mu_1} \dots \partial_{a_n} X^{\mu_n} \gamma_{\mu_1} \wedge \dots \wedge \gamma_{\mu_n} \quad (21)$$

After the integration over a finite n -surface Σ_B with boundary B we obtain a finite n -vector

$$\begin{aligned} \int_{\Sigma_B} d\Sigma &= \frac{1}{n!} X^{\mu_1 \dots \mu_n} \gamma_{\mu_1} \wedge \dots \wedge \gamma_{\mu_n} = \frac{1}{n!} \int_{\Sigma_B} d\xi^{a_1 \dots a_n} \partial_{a_1} X^{\mu_1} \dots \partial_{a_n} X^{\mu_n} \gamma_{\mu_1} \wedge \dots \wedge \gamma_{\mu_n} \\ &= \frac{1}{n!} \int_{\Sigma_B} d\xi^{a_1 \dots a_n} \frac{1}{n!} \partial_{[a_1} X^{\mu_1} \dots \partial_{a_n]} X^{\mu_n} \gamma_{\mu_1} \wedge \dots \wedge \gamma_{\mu_n} \end{aligned} \quad (22)$$

Its n -vector components are

$$X^{\mu_1 \dots \mu_n}[B] = \int_{\Sigma_B} d\xi^{a_1 \dots a_n} \frac{1}{n!} \partial_{[a_1} X^{\mu_1} \dots \partial_{a_n]} X^{\mu_n} \quad (23)$$

They describe an oriented n -area associated with Σ_B , whose boundary B will be called $(n-1)$ -loop, and $X^{\mu_1 \dots \mu_n}[B]$ are its *extrinsic coordinates*. With the same $(n-1)$ -loop we can associate *intrinsic coordinates* (parameters), in analogy to eqs. (13)–(16), according to

$$\xi^{a_1 \dots a_n}(B) = \int_{\Sigma_B} d\xi'^{a_1 \dots a_n} \quad (24)$$

With a particular choice of coordinates ξ^a , such that $\det \gamma_{ab} = 1$, the quantities $\xi^{a_1 \dots a_n}$ in the above equation determine the intrinsic (scalar) n -area of the n -surface bounded by the $(n-1)$ -loop.

As in the case of V_2 we assume that on our n -dimensional worldsheet V_n there exists a family of $(n-1)$ -loops B , described by functions $X^\mu(s^{\bar{a}}, \alpha)$, $\bar{a} = 1, 2, \dots, n-1$,

Instead of a manifold V_n of points we thus consider a family of loops. With every $(n-1)$ -loop of the family we can associate arbitrary parameters $\xi^{a_1 \dots a_n}$ (coordinates are like “house numbers”). Because of the property

$$\xi^{a_1 \dots a_j a_k \dots a_n} = -\xi^{a_1 \dots a_k a_j \dots a_n} \quad (25)$$

there is actually a single parameter. This is a particular choice for parameter α of our family of loops $X^\mu(s^{\bar{a}}, \alpha)$. By means of a mapping

$$\xi^{a_1 \dots a_n} \rightarrow x^{\mu_1 \dots \mu_n} = X^{\mu_1 \dots \mu_n}(\xi^{a_1 \dots a_n}) \quad (26)$$

we obtain a quenched minisuperspace description of a family of $(n-1)$ -loops, i.e., a description in terms of the target space multivector coordinate functions $X^{\mu_1 \dots \mu_n}$. The

family is such that the totality of the points of the $(n - 1)$ -loops belonging to the family is in one-to-one correspondence with the points of the worldsheet V_n . In other words, by mapping (26) we have *a quenched minisuperspace description of worldsheet*.

We started from a brane described by the embedding functions $X^\mu(\xi^a)$, and derived the expression (23) and functions (26). Once we have $X^{\mu_1 \dots \mu_2}$ as functions of a parameter $\xi^{a_1 \dots a_n}$, we may *forget* about the embedding $x^\mu = X^\mu(\xi^a)$ that we started from. *We may assume that all the information available to us are just functions $X^{\mu_1 \dots \mu_n}(\xi^{a_1 \dots a_n})$ given by mapping (26)*. Then we do not have knowledge of a particular worldsheet's manifold V_n , but of a class $\{V_n\}$ of worldsheet's manifolds that all satisfy eq. (26) for given functions $X^{\mu_1 \dots \mu_n}(\xi^{a_1 \dots a_n})$. So we calculate the derivative

$$\partial_{a_1 \dots a_n} X^{\mu_1 \dots \mu_n} \equiv \frac{\partial X^{\mu_1 \dots \mu_n}}{\partial \xi^{a_1 \dots a_n}} \quad (27)$$

which generalizes the notion of velocity.

In the above considerations one has to bear in mind that many loop configuration may cast the same holographic projections onto the coordinate planes as a single loop configuration. Therefore, a given set of polyvector coordinates $X^{\mu_1 \dots \mu_n}(\xi^{a_1 \dots a_n})$ may describe a single or many loop configuration. Not only the details of a single loop configuration (its infinite dimensionality), but also the number of loops is undetermined in this quenched description of loops.

3 The dynamics of extended objects

3.1 Objects described in terms of $X^\mu(\xi^a)$

The extended objects described by the mapping (1) obey the dynamical law that is incorporated in the Dirac-Nambu-Goto minimal surface action. An equivalent action that was considered in ref. [12] is a functional of the embedding functions $X^\mu(\xi^a)$ and the coordinate basis vector fields $e^a(\xi)$ having the role of Lagrange multipliers:

$$I[X^\mu, e^a] = \frac{\kappa}{2} \int d^n \xi |e| (e^a \partial_a X^\mu e^b \partial_b X_\mu + 2 - n) \quad (28)$$

where $|e| \equiv \sqrt{|\gamma|}$ is the determinant of $\gamma_{ab} = e_a \cdot e_b$.

Expanding the coordinate vector fields $e^a(\xi)$, $a = 1, 2, \dots, n$ in terms of orthonormal vector fields⁷ $e^{\mathbf{a}}$, $\mathbf{a} = 1, 2, \dots, n$, by means of a tetrad $e^{\mathbf{a}}_{\mathbf{a}}(\xi)$ according to

$$e^a(\xi) = e^{\mathbf{a}}_{\mathbf{a}}(\xi) e^{\mathbf{a}} \quad (29)$$

⁷Their inner products $e^{\mathbf{a}} \cdot e^{\mathbf{b}} = \eta^{\mathbf{ab}}$ gives the Minkowski metric.

we find the following relations⁸

$$\frac{\partial e^a}{\partial e^b} \equiv e_c \frac{\partial e^a}{\partial e^b_c} = n \delta^a_b \quad (30)$$

$$\frac{\partial |e|}{\partial e^a} = \frac{\partial |e|}{\partial \gamma^{cd}} \frac{\partial \gamma^{cd}}{\partial e^a} = -n |e| e_a \quad (31)$$

where n comes from the contraction $e^a e_a = n$.

Using (30),(31) we find that the variation of the action (28) with respect to e^a gives

$$-\frac{1}{2} e_c (e^a \partial_a X^\mu \partial^b X_\mu + 2 - n) + \partial_c X^\mu \partial_d X_\mu e^d = 0 \quad (32)$$

Performing the inner product with e^c and using $e^c \cdot e_c = n$ we find

$$e^a \partial_a X^\mu e^b \partial_b X_\mu = n \quad (33)$$

and eq.(32) becomes

$$e_c = \partial_c X^\mu \partial_d X_\mu e^d \quad (34)$$

This is the equation of “motion” for the Lagrange multipliers e_a . In order to understand better the meaning of eq.(34) let us perform the inner product with e_a :

$$e_c \cdot e_a = \partial_c X^\mu \partial_d X_\mu e^d \cdot e_a \quad (35)$$

Since $e_c \cdot e_a = \gamma_{ca}$ and $e^d \cdot e_a = \delta^d_a$ we obtain after renaming the indices

$$\gamma_{ab} = \partial_a X^\mu \partial_b X_\mu \quad (36)$$

This is the relation for the induced metric on the worldsheet. On the other hand, eq. (35) can be written as

$$e_a \cdot e_b = (\partial_a X^\mu \gamma_\mu) \cdot (\partial_a X^\nu \gamma_\nu) \quad (37)$$

from which we have that basis vectors e_a on the worldsheet V_n are expressed in terms of the embedding space basis vectors γ_μ :

$$e_a = \partial_a X^\mu \gamma_\mu \quad (38)$$

With our procedure we have thus derived eq.(4) as a solution to our dynamical sytem.

Using eq. (36) we find that the action (28) is equivalent to the well known Howe–Tucker action which is a functional of $X^\mu(\xi)$ and γ^{ab} :

$$I[X^\mu, \gamma_{ab}] = \frac{\kappa}{2} \int d^n \xi \sqrt{|\gamma|} (\gamma^{ab} \partial_a X^\mu \partial_b X_\mu + 2 - n) \quad (39)$$

⁸Eq.(30) also comes directly from the relation for a derivative with respect to a vector [9].

3.2 Objects described in terms of $X^{\mu_1 \dots \mu_n}(\xi^{a_1 \dots a_n})$

In Sec. 2.3 we have seen that an alternative description of extended objects, up to a class in which all objects have the same coordinates $X^{\mu_1 \dots \mu_n}$, is given by the mapping (26). Let us assume that such objects are described by the following action

$$I[X^{\mu_1, \dots, \mu_n}, e] = \frac{\kappa}{2} \int d^n \xi |e| \left[\frac{1}{n!} \left(\frac{1}{n!} e^{a_1} \wedge \dots \wedge e^{a_n} \frac{\partial X^{\mu_1 \dots \mu_n}}{\partial \xi^{a_1 \dots a_n}} \right)^\dagger \right. \\ \left. \times \left(\frac{1}{n!} e^{b_1} \wedge \dots \wedge e^{b_n} \frac{\partial X_{\mu_1 \dots \mu_n}}{\partial \xi^{b_1 \dots b_n}} \right) + 1 \right] \quad (40)$$

Factor $1/n!$ inside the bracket comes from the definition of the worldsheet n -vector $(1/n!)e^{a_1} \wedge \dots \wedge e^{a_n} \partial X^{\mu_1 \dots \mu_n} / \partial \xi^{a_1 \dots a_n}$. The extra factor $1/n!$ in front of the bracket comes from the square of the target space n -vector $(1/n!)(\partial X^{\mu_1 \dots \mu_n} / \partial \xi^{a_1 \dots a_n}) \gamma_{\mu_1} \wedge \dots \wedge \gamma_{\mu_n}$. The operation $\dagger$ reverses the order of vectors.

Let us take into account the following relations:

$$e_{a_1} \wedge \dots \wedge e_{a_n} = e \epsilon_{a_1 \dots a_n} \quad (41)$$

$$e^{a_1} \wedge \dots \wedge e^{a_n} = e^{-1} \epsilon^{a_1 \dots a_n} \quad (42)$$

$$e^{-1} = \frac{e}{|e|^2}, \quad |e| \equiv \sqrt{e^\dagger \cdot e} = \sqrt{|\gamma|} \equiv \lambda \quad (43)$$

$$\gamma = \det \gamma_{ab}, \quad \gamma_{ab} = e_a \cdot e_b \quad (44)$$

Instead of the intrinsic parameters $\xi^{a_1 \dots a_n}$, let us introduce the dual parameter

$$\tilde{\xi} = \frac{1}{n!} \epsilon_{a_1 \dots a_n} \xi^{a_1 \dots a_n} \quad (45)$$

and rewrite the n -area velocity according to

$$\frac{\partial X^{\mu_1 \dots \mu_n}}{\partial \xi^{c_1 \dots c_n}} = \frac{\partial X^{\mu_1 \dots \mu_n}}{\partial \tilde{\xi}} \frac{\partial \tilde{\xi}}{\partial \xi^{c_1 \dots c_n}} = \frac{\partial X^{\mu_1 \dots \mu_n}}{\partial \tilde{\xi}} \epsilon_{c_1 \dots c_n} \quad (46)$$

where we have used

$$\frac{\partial \xi^{a_1 \dots a_n}}{\partial \xi^{c_1 \dots c_n}} = \delta_{c_1 \dots c_n}^{a_1 \dots a_n} \quad (47)$$

and where the generalized Kronecker symbol is given by the antisymmetrized sum of products of ordinary deltas. From eq.(46) we have

$$\frac{\partial X^{\mu_1 \dots \mu_n}}{\partial \tilde{\xi}} = \frac{1}{n!} \epsilon^{c_1 \dots c_n} \frac{\partial X^{\mu_1 \dots \mu_n}}{\partial \xi^{c_1 \dots c_n}} \quad (48)$$

By using eqs.((33),(35) and (48) we can rewrite the action (40) as

$$I[X^{\mu_1 \dots \mu_n}, \lambda] = \frac{\kappa}{2} \int d\tilde{\xi} \left(\frac{1}{\lambda n!} \frac{\partial X^{\mu_1 \dots \mu_n}}{\partial \tilde{\xi}} \frac{\partial X_{\mu_1 \dots \mu_n}}{\partial \tilde{\xi}} + \lambda \right) \quad (49)$$

where

$$d\tilde{\xi} = \frac{1}{n!} \epsilon_{a_1 \dots a_n} d\xi^{a_1 \dots a_n} = \frac{1}{n!} \epsilon_{a_1 \dots a_n} d\xi_a^{[a_1} \dots d\xi_n^{a_n]} = \epsilon_{a_1 \dots a_n} d\xi_a^{a_1} \dots d\xi_n^{a_n} = d\xi^1 d\xi^2 \dots d\xi^n \equiv d^n \xi \quad (50)$$

The last step in eq.(50) holds in a coordinates system in which $d\xi_1^{a_1} = d\xi^1$, $d\xi_2^{a_2} = d\xi^2$, ..., $d\xi_n^{a_n} = d\xi^n$.

The action (49) is a functional of the n -area variables $X^{\mu_1 \dots \mu_n}(\tilde{\xi})$ and a Lagrange multiplier λ , defined in eq.(43). Variation of eq. (49) with respect to λ and $X^{\mu_1 \dots \mu_n}$, respectively, gives

$$\delta\lambda : \quad \frac{1}{n!} \frac{\partial X^{\mu_1 \dots \mu_n}}{\partial \tilde{\xi}} \frac{\partial X_{\mu_1 \dots \mu_n}}{\partial \tilde{\xi}} - \lambda^2 = 0 \quad (51)$$

$$\delta X^{\mu_1 \dots \mu_n} : \quad \frac{d}{d\tilde{\xi}} \left(\frac{1}{\lambda} \frac{\partial X^{\mu_1 \dots \mu_n}}{\partial \tilde{\xi}} \right) = 0 \quad (52)$$

These are the equations of motion for the n -area variables.

Inserting eq.(51) into (49) we obtain the action which is a functional of $X^{\mu_1 \dots \mu_n}$ solely:

$$I[X^{\mu_1 \dots \mu_n}] = \kappa \int d\tilde{\xi} \left(\frac{1}{n!} \frac{\partial X^{\mu_1 \dots \mu_n}}{\partial \tilde{\xi}} \frac{\partial X_{\mu_1 \dots \mu_n}}{\partial \tilde{\xi}} \right)^{1/2} \quad (53)$$

The latter action has the same form as the action for a worldline of a relativistic particle. Factor $1/n!$ in the latter action disappears, if we order indices according to $\mu_1 < \mu_2 < \dots < \mu_n$.

3.3 More general objects (generalization to C -space)

So far we have considered branes described by coordinate functions of the type $X^\mu(\xi)$, $X^{\mu_1 \mu_2}(\xi^{a_1 a_2})$, ..., or $X^{\mu_1 \dots \mu_n}(\xi^{a_1 \dots a_n})$ that represent a mapping of a worldsheet loop into a target space loop of the same dimensionality.:

$$(n-1)\text{-loop on } V_n \longrightarrow (n-1)\text{-loop in } V_N.$$

Let us now extend the theory and consider “mixed” mappings

$$(r-1)\text{-loop on } V_n \longrightarrow (R-1)\text{-loop in } V_N.$$

where r , in general, is different from R . So we arrive at a more general extended object which is described by a mapping [12, 15]

$$\xi^{a_1 \dots a_r} \rightarrow x^{\mu_1 \dots \mu_R} = X^{\mu_1 \dots \mu_R}(\xi, \xi^a, \xi^{a_1 a_2}, \dots, \xi^{a_1 \dots a_s}, \dots, \xi^{a_1 \dots a_n}), \quad 0 \leq R \leq N, \quad 0 \leq r \leq n < N \quad (54)$$

In the compact notation we set

$$\begin{aligned} X^M &\equiv X^{\mu_1 \dots \mu_R}, & \mu_1 < \mu_2 < \dots < \mu_R \\ \xi^A &\equiv \xi^{a_1 \dots a_r}, & a_1 < a_2 < \dots < a_r \end{aligned}$$

and write the mapping (54) as

$$\xi^A \rightarrow x^M = X^M(\xi^A) \quad (55)$$

This is the parametric equation of our generalized extended object. Such object lives in a target space which is now generalized to *Clifford space* (shortly *C-space*). The worldsheet associated with the extended object is also generalized to a Clifford space. In the following we will explain this in more detail.

In eq. (54) or (55) we have a generalization of the usual relation

$$\xi^a \rightarrow x^\mu = X^\mu(\xi^a), \quad a = 1, 2, \dots, n; \quad \mu = 0, 1, 2, \dots, N-1 \quad (56)$$

that describes an n -dimensional surface, called worldsheet or world manifold, V_n , embedded in N -dimensional target space V_N . In eq. (56) the space $\mathbb{R}^n$ of parameters ξ^a is isomorphic to an n -dimensional vector space $\mathbf{V}_n$, spanned by an orthonormal basis $\{h_a\}$. The *vector space* $\mathbf{V}_n$ should not be confused with the worldsheet V_n , which is a *manifold* (embedded in a higher dimensional manifold V_N).

Instead of $\mathbf{V}_n$ we can consider the corresponding Clifford algebra $\mathbf{C}_n$ which is itself a vector space. Amongst its elements are r -vectors associated with $(r-1)$ -loops, $r = 0, 1, 2, \dots, n$. A generic object is a superposition of r -vectors for different grades, and it is described by a Clifford number, a polyvector, $\xi^{a_1 \dots a_r} h_{a_1} \wedge \dots \wedge h_{a_r} \in \mathbf{C}_n$.

Our objects are now *extended events* $\mathcal{E}$ [29], superpositions of $(r-1)$ -loops, to which we assign a set of 2^n parameters (coordinates) $\xi^A \equiv \xi^{a_1 \dots a_r}$, $r = 0, 1, 2, \dots, n$ according to the mapping

$$\mathcal{E} \rightarrow \xi^A(\mathcal{E}) \quad (57)$$

The assignment is arbitrary. We may choose an object $\mathcal{E}_0$ to which we assign coordinates $\xi^A(\mathcal{E}) = 0$. This is a coordinate origin. Choosing an origin $\mathcal{E}_0$, the polyvectors $\xi^A h_A$ pointing from $\mathcal{E}_0$ to any $\mathcal{E}$ are in one-to-one correspondence with extended events $\mathcal{E}$. The space of extended events is then isomorphic to Clifford algebra $\mathbf{C}_n$, and the latter algebra, in turn, is isomorphic to the space of parameter $\{\xi^A\} = \mathbb{R}^{2^n}$. Therefore we will speak about $\mathbf{C}_n$ as the *parametric Clifford algebra* or *parametric polyvector space*.

The parametric space $\mathbf{C}_n$ is by definition a (*poly*)*vector space*, spanned by a basis $h_{a_1} \wedge \dots \wedge h_{a_r}$, $r = 0, 1, 2, \dots, n$, formed by the *orthonormal basis* $\{h_a\}$, $a = 1, 2, \dots, n$. This implies that $\mathbf{C}_n$ is a metric space, but its metric is just formal, without any physical content. Now let us consider the mapping (55) from $\mathbf{C}_n$ into a Clifford space C_{V_N} generated by a

basis γ_μ , $\mu = 0, 1, 2, \dots, N - 1$, with $N > n$. So we obtain a generalized, 2^n -dimensional surface C_{V_n} embedded in a target Clifford space C_{V_N} . The surface C_{V_n} generalizes the notion of worldsheet V_n to Clifford space, i.e., a manifold such that any of its tangents spaces is a Clifford algebra. If we consider only the intrinsic properties of C_{V_n} (i.e., if we “forget” about its embedding into a higher dimensional Clifford space C_{V_N}), then we can simply denote it as C_n .

Bellow we summarize our notation of various spaces:

- $\mathbf{V}_n$ *Parametric vector space*, with an orthonormal basis $\{h_a\}$, $a = 1, 2, \dots, n$ and elements $\xi^a h_a \in \mathbf{V}_n$. It is isomorphic to $\mathbb{R}^n$, the space of parameters ξ^a .
- V_n *Manifold*, either flat or curved. It is a space of points (events) $\mathcal{P}$. With every point $\mathcal{P} \in V_n$ we associate a set of n parameters (coordinates) $\xi^a(\mathcal{P}) \equiv \xi^a \in \mathbb{R}^n$. Coordinate basis vectors are e^a , whilst orthonormal basis vectors are $e_{\mathbf{a}}$.
- $\mathbf{C}_n$ *Parametric Clifford algebra* of $\mathbf{V}_n$, called also *parametric polyvector space*, with basis $\{h_A\} \equiv \{h_{a_1} \wedge \dots \wedge h_{a_r}\}$, $r = 0, 1, 2, \dots, n$ and elements $\xi^A h_A \in \mathbf{C}_n$, called *polyvectors*. It is isomorphic to $\mathbb{R}^{2^n}$, the space of parameters ξ^A .
- C_n *Clifford manifold*, or *Clifford space*, either flat or curved. It is a space of points that are perceived in a subspace V_n as *extended events* $\mathcal{E}$. With every $\mathcal{E} \in C_n$ we associate a set of 2^n parameters $\xi^A(\mathcal{E}) \equiv \xi^A \equiv \xi^{a_1 \dots a_r} \in \mathbb{R}^{2^n}$, $r = 0, 1, 2, \dots, n$. Coordinate basis elements are $e_A \equiv e_{a_1 \dots a_r}$; orthonormal basis elements are $e_{\mathbf{A}} \equiv e_{\mathbf{a}_1 \dots \mathbf{a}_r}$, $r = 0, 1, 2, \dots, n$. In particular C_n can be considered as being embedded in a higher dimensional Clifford space; then it is denoted as C_{V_n} .
- C_{V_n} *Generalized worldsheet*, a Clifford space embedded in a target Clifford space. Its coordinate basis elements are $e^A \equiv e_{a_1 \dots a_r}$; orthonormal basis elements are $e_{\mathbf{A}} \equiv e_{\mathbf{a}_1 \dots \mathbf{a}_r}$, $r = 0, 1, 2, \dots, n$.
- C_{V_N} *Target Clifford manifold*, or *target Clifford space*, either flat or curved. It is a space of points that are perceived in a subspace V_N as extended events $\mathcal{E}$. With every $\mathcal{E} \in C_{V_N}$ we associate a set of 2^N coordinates $x^M(\mathcal{E}) \equiv x^M \equiv x^{\mu_1 \dots \mu_R}$, $R = 0, 1, 2, \dots, N$. Coordinate basis elements are $\gamma_M \equiv \gamma_{\mu_1 \dots \mu_R}$; orthonormal basis elements are $\gamma_{\mathbf{M}} \equiv \gamma_{\mu_1 \dots \mu_R}$. Instead of C_{V_N} we can use simply notation C_N .

We follow the rule that bold symbols are used for vector spaces, whilst light symbols are used for manifolds. Since Clifford algebras also are vector spaces, they are denoted by bold symbols, whereas the corresponding Clifford spaces (manifolds of points representing extended events) are denoted by light symbols. By such notation we have attempted to simplify distinction among all those various spaces that occur in our theory of generalized

branes. For these purely physical reasons we have thus, to certain extent, deviated from the standard notation used in mathematics.

An infinitesimal (polyvector) $dX \in C_{V_N}$, joining two points on the surface C_{V_n} can be written as

$$dX = dX^M \gamma_M = d\xi^A \partial_A X^M \gamma_M = d\xi^A e_A \quad (58)$$

where

$$e_A = \partial_A X^M \gamma_M \quad (59)$$

These are induced basis tangent (polyvectors) on C_{V_n} .

At every point of the *flat* target C -space C_{V_N} there exists a basis

$$\{\gamma^M\} = \{\mathbf{1}, \gamma^\mu, \gamma^{\mu_1\mu_2}, \dots, \gamma^{\mu_1\dots\mu_N}\} \quad (60)$$

given in terms of 2^N Clifford numbers

$$\gamma^M \equiv \gamma^{\mu_1\dots\mu_R} \equiv \gamma^{\mu_1} \wedge \gamma^{\mu_2} \wedge \dots \wedge \gamma^{\mu_R}, \quad \mu_1 < \mu_2 < \dots \mu_R, \quad R = 0, 1, 2, \dots, N \quad (61)$$

At every point $\xi \in C_{V_n}$ on the brane's worldsheet C -space C_{V_n} , which in general is *curved*, there exist a basis given in terms of 2^n Clifford numbers⁹

$$\{e^A\} = \{e, e^a, e^{a_1a_2}, \dots, e^{a_1\dots a_n}\} \quad (62)$$

that span a tangent space $T_\xi(C_{V_n})$. At a particular point $\xi_0 \in C_{V_n}$ it may hold

$$e^A \equiv e^{a_1\dots a_r} \equiv e^{a_1} \wedge e^{a_2} \wedge \dots \wedge e^{a_r}, \quad a_1 < a_2 < \dots < a_r, \quad r = 0, 1, 2, \dots, n \quad (63)$$

That is, at that particular point the basis polyvectors on C_{V_n} are given as wedge products of basis vectors e^a . The above property (63) cannot hold globally on a *curved* C_{V_n} . At points different from ξ_0 , basis polyvectors are in general superpositions of $e^{a_1} \wedge e^{a_2} \wedge \dots \wedge e^{a_s}$, $s = 0, 1, 2, \dots, n$.

To sum up, every Clifford number in the target C -space can be expanded in terms of γ^M , and every Clifford number on the worldsheet C -space can be expanded in terms of e^A . Such numbers are also called *Clifford aggregates* or *polyvectors*. They are superpositions r -vectors, the objects of definite grade that we call multivectors. This description automatically contains *spinors* which are just members of the left or right ideal of Clifford algebra [27].

⁹We will use ‘basis’ and ‘frame’ as synonyms. In order to simplify notation and wording we will be sloppy in distinguishing objects from the corresponding fields, e.g., (poly)vectors from (poly)vector fields, frames from frame fields, etc. From the context it should not be difficult to understand which ones we have in mind.

Metric of C_{V_N} is $G_{MN} = \gamma_M^\dagger * \gamma_N$, whilst metric of C_{V_n} is $G_{AB} = e_A^\dagger * e_B$. Here ‘ $*$ ’ denotes the scalar product of two Clifford numbers A and B

$$A * B = \langle AB \rangle_0 \quad (64)$$

Let us now define the object V which is a polyvector in target space and on the worldsheet:

$$V = e^A \frac{\partial X^M}{\partial \xi^A} \gamma_M = \sum_{r=0}^N \sum_{s=0}^n \frac{1}{r!s!} e^{a_1 \dots a_s} \frac{\partial X^{\mu_1 \dots \mu_r}}{\partial \xi^{a_1 \dots a_s}} \gamma_{\mu_1 \dots \mu_r} \quad (65)$$

In the right hand side expression we impose no restriction on the indices $\mu_1, \mu_2, \dots, \mu_r$ and $a_1, a_2, \dots, a_s$, therefore, in order to prevent multiple counting of equivalent terms, factors $1/r!$ and $1/s!$ are introduced.

It is illustrative to express eq.(65) in a more explicit form by employing the notation (60)–(63) and by writing

$$X^M = (\Omega, X^\mu, X^{\mu_1 \mu_2}, \dots, X^{\mu_1 \dots \mu_N}) \quad (66)$$

$$\xi^A = (\xi, \xi^a, \xi^{a_1 a_2}, \dots, \xi^{a_1 \dots a_n}) \quad (67)$$

We obtain

$$\begin{aligned} V = & \left(\frac{\partial \Omega}{\partial \xi} + e^a \frac{\partial \Omega}{\partial \xi^a} + \frac{1}{2!} e^{a_1 a_2} \frac{\partial \Omega}{\partial \xi^{a_1 a_2}} + \dots \frac{1}{n!} e^{a_1 \dots a_n} \frac{\partial \Omega}{\partial \xi^{a_1 \dots a_n}} \right) \mathbb{1} \\ & + \left(\frac{\partial X^\mu}{\partial \xi} + e^a \frac{\partial X^\mu}{\partial \xi^a} + \frac{1}{2!} e^{a_1 a_2} \frac{\partial X^\mu}{\partial \xi^{a_1 a_2}} + \dots \frac{1}{n!} e^{a_1 \dots a_n} \frac{\partial X^\mu}{\partial \xi^{a_1 \dots a_n}} \right) \gamma_\mu \\ & + \frac{1}{2!} \left(\frac{\partial X^{\mu_1 \mu_2}}{\partial \xi} + e^a \frac{\partial X^{\mu_1 \mu_2}}{\partial \xi^a} + \frac{1}{2!} e^{a_1 a_2} \frac{\partial X^{\mu_1 \mu_2}}{\partial \xi^{a_1 a_2}} + \dots \frac{1}{n!} e^{a_1 \dots a_n} \frac{\partial X^{\mu_1 \mu_2}}{\partial \xi^{a_1 \dots a_n}} \right) \gamma_{\mu_1 \mu_2} \\ & + \\ & \vdots \\ & + \frac{1}{N!} \left(\frac{\partial X^{\mu_1 \dots \mu_N}}{\partial \xi} + e^a \frac{\partial X^{\mu_1 \dots \mu_N}}{\partial \xi^a} + \frac{1}{2!} e^{a_1 a_2} \frac{\partial X^{\mu_1 \dots \mu_N}}{\partial \xi^{a_1 a_2}} + \dots \frac{1}{n!} e^{a_1 \dots a_n} \frac{\partial X^{\mu_1 \dots \mu_N}}{\partial \xi^{a_1 \dots a_n}} \right) \gamma_{\mu_1 \dots \mu_N} \end{aligned} \quad (68)$$

One might ask how such a generalized extended object, described by eq.(55), that sweep a Clifford worldsheet C_{V_n} , embedded in a Clifford space C_{V_N} , looks like. Here our perceptive system again shows its shortcomings, like in the case of figuring out how higher dimensional objects look like. We are able to draw pictures of projections of an object onto 3 or 2-dimensional Euclidean space, and that is, more or less, all. But on the other hand, we are able to do algebra, and the algebra is interpreted as geometric algebra. So we have to content us by our ability to control the situation algebraically, and assume that there is a mapping between algebraic and geometric objects. The latter objects are associated with physical objects, such as, e.g., the generalized extended objects, that incorporate branes of various dimensionalities.

Let the action describing the dynamics of a generalized extended object, shortly, a generalized brane, be described by the embedding functions $X^M(\xi^A)$ be

$$I[X^M, e^A] = \frac{T}{2} \int d^{d_C} \xi |E| \left[\left(e^A \frac{\partial X^M}{\partial \xi^A} \gamma_M \right)^\dagger * \left(e^B \frac{\partial X^N}{\partial \xi^B} \gamma_N \right) + 2 - d_C \right] \quad (69)$$

The latter action has a similar form as the action (28). But, since the indices M, N , and A, B run over the full basis (60) and (62) of the corresponding C -spaces, the action (69) is more general than (28).

Here the measure $d^{d_C} \xi |E|$ is the volume element in the worldsheet C -space C_{V_n} (whose dimension is 2^{d_C}). It is equal to the product

$$d^{d_C} \xi |E| \equiv |E| \prod_{A_i} d\xi^{A_i} = d\xi \prod_{a_1} d\xi^{a_1} \prod_{a_1 < a_2} d\xi^{a_1 a_2} \dots \prod_{a_1 < \dots < a_n} d\xi^{a_1 \dots a_n} |E| \quad (70)$$

By $|E|$ we denote the square root of the determinant of the worldsheet C -space metric which is given by the scalar product

$$G^{AB} = (e^A)^\dagger * e^B = \langle (e^A)^\dagger e^B \rangle_0 \quad (71)$$

where $\langle \rangle_0$ denotes the scalar part. Explicitly,

$$|E| = \sqrt{|G|}, \quad G \equiv \det G_{AB} = \frac{1}{d_C!} \epsilon^{A_1 \dots A_{d_C}} \epsilon^{B_1 \dots B_{d_C}} G_{A_1 B_1} \dots G_{A_{d_C} B_{d_C}} \quad (72)$$

The action (69) is a functional of X^M and e^A , and is a C -space generalization of the action (28) which is a functional of the worldsheet embedding functions X^μ and basis vectors e^a .

An action which is classically equivalent to (69) is a functional of X^M and G^{AB} :

$$I[X^M, G^{AB}] = \frac{T}{2} \int d^{d_C} \xi \sqrt{|G|} (G^{AB} \partial_A X^M \partial_B X_M + 2 - d_C) \quad (73)$$

where $\partial_B X_M = G_{MN} \partial_B X^N$, $G_{MN} = \gamma_M^\dagger * \gamma_N$. In eq.(73) we have a C -space generalization of the well known Howe–Tucker action [34].

Variation of the action (73) with respect to X^M and G^{AB} gives the equations of motion of our C -space extended object:

$$\delta X^M : \quad \frac{1}{\sqrt{|G|}} \partial_A (\sqrt{|G|} \partial^A X^M) = 0 \quad (74)$$

$$\delta G^{AB} : \quad G_{AB} = \partial_A X^M \partial_B X_M \quad (75)$$

Taking into account that $G_{AB} G^{AB} = d_C = 2^n$ and inserting eq. (75) Into eq. (73) we obtain the action integral

$$I[X^M] = T \int d^{d_C} \xi \sqrt{\det \partial_A X^M \partial_B X_M} \quad (76)$$

which is the volume of the C -space worldsheet. The latter action contains the usual p -branes, including point particles, as special cases.

Our action (73), or equivalently (76), is invariant under reparametrizations of coordinates ξ^A , $A = 1, 2, \dots, 2^n$. As a consequence, there are 2^n primary constraints. So we are free to choose 2^n relations among our dynamical variables $X^M(\xi^A)$, and thus fix a gauge (a parametrization). For one of those relations we can choose, for instance,

$$\partial_{a_1 \dots a_r} X^{\mu_1 \dots \mu_r} = \partial_{[a_1} X^{\mu_1} \dots \partial_{a_r]} X^{\mu_r} \quad (77)$$

That is, the above relation is just one of gauge fixing relations. It will be used in Sec. 5.3, where we will consider the coupling of our generalized branes to external fields.

4 On the relativity in Clifford space

The discussion of previous sections has led us to the conclusion that the space in which our extended objects live is *Clifford space*, shortly C -space, denoted C_{V_N} or C_N . A point in C -space is described by the coordinates $x^M = (\Omega, x^\mu, x^{\mu_1 \mu_2}, \dots, x^{\mu_1 \dots \mu_N})$ which together with the basis elements $\gamma^M = (1, \gamma_\mu, \gamma_{\mu_1 \mu_2}, \dots, \gamma_{\mu_1 \dots \mu_N})$ form a coordinate polyvector¹⁰

$$X = x^M \gamma_M = \frac{1}{r!} \sum_{r=0}^N x^{\mu_1 \dots \mu_r} \gamma_{\mu_1 \dots \mu_r} \quad (78)$$

From the point of view of 2^N -dimensional C -space, $x^{\mu_1 \dots \mu_r}$, $r = 1, \dots, N$, are coordinates of a point, whilst from the point of view of the underlying Minkowski space V_N , these are the r -area (r -volume) variables.

The infinitesimal polyvector connecting two infinitesimally separated points of C -space is

$$dX = dx^M \gamma_M = \frac{1}{r!} \sum_{r=0}^N dx^{\mu_1 \dots \mu_r} \gamma_{\mu_1 \dots \mu_r} \equiv dx^M \gamma_M \quad (79)$$

The square of the distance between these points is given by the scalar product

$$|dX|^2 \equiv dX^\dagger * dX = dx^M dx^N G_{MN} = dx^M dx_N \quad (80)$$

where G_{MN} is the metric of C -space¹¹ :

$$G_{MN} = \gamma_M^\dagger * \gamma_N = \gamma_{\mu_1 \dots \mu_r}^\dagger * \gamma_{\nu_1 \dots \nu_s} \quad (81)$$

¹⁰ In flat C -space it makes sense to consider a polyvector joining the coordinate origin $\mathcal{E}_0$ with coordinates $x^M(\mathcal{E}_0) = 0$ and a point $\mathcal{E}$ with coordinates $x^M(\mathcal{E}) \equiv x^M$, where $\mathcal{E}_0, \mathcal{E} \in C_{V_N}$.

¹¹ A reason why we define the C -space metric by employing the reversion is in the consistency between G_{MN} , its inverse G^{MN} , and the relations (82), (83) (in which the indices $\mu_i, \nu_j, \dots$ are lowered and raised by the ordinary 4-dimensional metric $g_{\mu_i \nu_j}$ and its inverse $g^{\mu_i \nu_j}$). For more details see ref. [16].

In particular, C -space C_{V_N} can be flat. This means that the curvature of the connection on flat C_{V_N} vanishes (see sec. 5.1). In such a case one can find a coordinates system on C_{V_N} such that the metric G_{MN} is diagonal.

Explicitly, for different choices of the indices M and N we have:

$$\begin{aligned}
G_{\mu\nu} &= \gamma_\mu \cdot \gamma_\nu = g_{\mu\nu} = \eta_{\mu\nu} \\
G_{[\mu_1\mu_2][\nu_1\nu_2]} &= \gamma_{\mu_1\mu_2}^\dagger * \gamma_{\nu_1\nu_2} = (\gamma_{\mu_2} \wedge \gamma_{\mu_1}) \cdot (\gamma_{\nu_1} \wedge \gamma_{\nu_2}) = \begin{vmatrix} g_{\mu_1\nu_1} & g_{\mu_1\nu_2} \\ g_{\mu_2\nu_1} & g_{\mu_2\nu_2} \end{vmatrix} \\
G_{[\mu_1\mu_2\mu_3][\nu_1\nu_2\nu_3]} &= \gamma_{\mu_1\mu_2\mu_3}^\dagger * \gamma_{\nu_1\nu_2\nu_3} = (\gamma_{\mu_3} \wedge \gamma_{\mu_2} \wedge \gamma_{\mu_1}) \cdot (\gamma_{\nu_1} \wedge \gamma_{\nu_2} \wedge \gamma_{\nu_3}) = \begin{vmatrix} g_{\mu_1\nu_1} & g_{\mu_1\nu_2} & g_{\mu_1\nu_3} \\ g_{\mu_2\nu_1} & g_{\mu_2\nu_2} & g_{\mu_2\nu_3} \\ g_{\mu_3\nu_1} & g_{\mu_3\nu_2} & g_{\mu_3\nu_3} \end{vmatrix} \\
G_{\mu[\nu_1\nu_2]} &= \gamma_\mu * \gamma_{\nu_1\nu_2} = \langle \gamma_\mu (\gamma_{\nu_1} \wedge \gamma_{\nu_2}) \rangle_0 = 0
\end{aligned} \tag{82}$$

In general we have

$$G_{[\mu_1\ldots\mu_r][\nu_1\ldots\nu_r]} = (\gamma_{\mu_r} \wedge \ldots \wedge \gamma_{\mu_1}) \cdot (\gamma_{\nu_1} \wedge \ldots \wedge \gamma_{\nu_r}) = \det g_{\mu_i\nu_j}, \quad i, j = 1, 2, \dots, r \tag{83}$$

when $r = s$, and

$$G_{[\mu_1\ldots\mu_r][\nu_1\ldots\nu_s]} = 0 \tag{84}$$

when $r \neq s$.

Taking into account the explicit expressions for metric (82)–(84), the quadratic form (80) reads

$$\begin{aligned}
|dX|^2 &= \frac{1}{r!} \sum_{r=0}^N dx^{\mu_1\ldots\mu_r} dx_{\mu_1\ldots\mu_r} \\
&= d\Omega^2 + dx^\mu dx_\mu + \frac{1}{2!} dx^{\mu_1\mu_2} dx_{\mu_1\mu_2} + \ldots + \frac{1}{N!} dx^{\mu_1\ldots\mu_N} dx_{\mu_1\ldots\mu_N}
\end{aligned} \tag{85}$$

In the latter expression only the factor $1/r!$ remains, since the other factor is canceled by $r!$ coming from the determinant (83). Indices $\mu_1, \mu_2, \dots$ are lowered and raised by Minkowski metric $\eta_{\mu\nu}$ and its inverse $\eta^{\mu\nu}$.

In the 16-dimensional Clifford space C_{M_4} of the 4-dimensional Minkowski spacetime the polyvector (79) and its square (80) can be rewritten as

$$dX = d\Omega + dx^\mu \gamma_\mu + \frac{1}{2} dx^{\mu\nu} \gamma_\mu \wedge \gamma_\nu + d\tilde{x}^\mu \gamma_5 \gamma_\mu + d\tilde{\Omega} \gamma_5 \tag{86}$$

$$|dX|^2 = d\Omega^2 + dx^\mu dx_\mu + \frac{1}{2} dx^{\mu\nu} dx_{\mu\nu} - d\tilde{x}^\mu \tilde{x}_\mu - d\tilde{s}^2 \tag{87}$$

where

$$d\tilde{x}^\mu \equiv \frac{1}{3!} dx^{\alpha\beta\rho} \epsilon_{\alpha\beta\rho}{}^\mu \tag{88}$$

$$d\tilde{\Omega} \equiv \frac{1}{4!} dx^{\alpha\beta\rho\sigma} \epsilon_{\alpha\beta\rho\sigma} \quad (89)$$

The minus sign in the last two terms of the above quadratic form occurs because in 4-dimensional spacetime with signature $(+---)$ we have $\gamma_5^2 = (\gamma_0\gamma_1\gamma_2\gamma_3)(\gamma_0\gamma_1\gamma_2\gamma_3) = -1$, and also $\gamma_5^\dagger\gamma_5 = (\gamma_3\gamma_2\gamma_1\gamma_0)(\gamma_0\gamma_1\gamma_2\gamma_3) = -1$.

It is illustrative to write the quadratic form (line element) explicitly:

$$\begin{aligned} |dX|^2 = & d\Omega^2 + (dx^0)^2 - (dx^1)^2 - (dx^2)^2 - (dx^3)^2 \\ & - (dx^{01})^2 - (dx^{02})^2 - (dx^{03})^2 + (dx^{12})^2 + (dx^{13})^2 + (dx^{23})^2 \\ & - (d\tilde{x}^0)^2 + (d\tilde{x}^1)^2 + (d\tilde{x}^2)^2 + (d\tilde{x}^3)^2 - d\tilde{\Omega}^2 \end{aligned} \quad (90)$$

Here $\tilde{x}^0 = x^{123}$, $\tilde{x}^1 = x^{023}$, $\tilde{x}^2 = x^{013}$, $\tilde{x}^3 = x^{012}$, $\tilde{\Omega} = x^{0123}$. The factor $1/2$ has disappeared from the term $(1/2)dx^{\mu\nu}dx_{\mu\nu}$, since $(1/2)(dx^{01}x_{01} + dx^{10}x_{10}) = dx^{01}dx_{01} = -(dx^{01})^2$, etc..

By inspecting the quadratic form (87) we see that it has 8 terms with *plus* and 8 terms with *minus* sign. The group of transformations that leave the quadratic form (87) invariant is $SO(8,8)$. These are pseudorotations in C -space and have the role of generalized Lorentz transformations:

$$x'^M = L^M_N x^N \quad (91)$$

The transformations matrix satisfies the relation $L^M_J L^N_K G_{MN} = G_{JK}$.

If interpreted actively (91) are the transformations that transform a point of C -space with coordinates x^M into another point with coordinates x'^M . In the passive interpretation the point remains the same, but its coordinates x^M with respect to a reference frame S are transformed into coordinates x'^M with respect to a reference frame S' .

From the point of view of C -space, x^M are coordinates of a *point*. But from the point of view of the underlying Minkowski space, x^M are the $(p+1)$ -vector (holographic) coordinates of p -loops associated with an extended object [11]. In C -space, p -loops of different dimensionalities p (i.e., points, closed lines, closed 2-surfaces, closed 3-volumes) are all described on the same footing [36, 37, 26, 11, 14], and can be transformed into each other by transformations (91). Pseudo rotations in C -space have thus the role of polydimensional rotations in M_4 . A point can be transformed into a line, and in general an $(R-1)$ -loop into an $(R'-1)$ -loop. So the very dimensionality of a loop can change under a transformation. This means that, when observed in a given reference frame S , a loop's dimensionality can change from $(R-1)$ to $(R'-1)$. Alternatively, dimensionality of the same loop, when observed from different reference frames S and S' , can look different. In short, dimensionality of a loop depends on the observer (associated with a given reference frame). Such relativity of dimensionality of a loop also explains why in the mapping (54),(55) the dimensionality of a loop in the parameter space $\{\xi^a\} \equiv \mathbb{R}^n$ is in general

different from the dimensionality of the same loop with respect to a reference frame in the target space V_N .

C -space in essence encodes the zero modes of p -loop configurations, since p -loops space is *infinite* dimensional whereas C -space is finite dimensional. As already mentioned before, a p -loop configuration, in general, can consist of many loops.

The construction with C -space coordinates, $SO(8,8)$ symmetry and the brane equations of motion (74) reminds us of the constructions considered in refs. [5], where extra coordinates were introduced in order to make manifest the $SO(n,n)$ symmetry of the duality transformations for strings and branes.

5 Curved Clifford space

5.1 General considerations

In general, the worldsheet V_n swept by a brane is *curved*. In Sec. 3.2 we have considered a concept of a more complicated, generalized brane, whose (generalized) worldsheet C_{V_n} was *curved Clifford space*. The latter worldsheet C_{V_n} was embedded in a target space which was a *flat Clifford space* C_{V_N} , with the metric properties given in Sec. 4.

A next step is to consider *curved target Clifford space* C_{V_N} (see refs. [28, 29]). At every point $\mathcal{E} \in C_{V_N}$ we have a flat tangent Clifford space $T_{\mathcal{E}}(C_{V_N})$ and an orthonormal basis of 2^N Clifford numbers

$$\{\gamma_{\mathbf{M}}\} = \{\mathbf{1}, \gamma_{\mu_1}, \gamma_{\mu_1\mu_2}, \dots, \gamma_{\mu_1\mu_2\dots\mu_N}\} \quad (92)$$

where

$$\gamma_{\mu_1\mu_2\dots\mu_r} = \gamma_{\mu_1} \wedge \gamma_{\mu_2} \wedge \dots \wedge \gamma_{\mu_r} \quad (93)$$

From an *orthonormal basis* $\{\gamma_{\mathbf{M}}\}$ we can switch to a *coordinate basis*

$$\{\gamma_M\} = \{\gamma, \gamma_{\mu_1}, \gamma_{\mu_1\mu_2}, \dots, \gamma_{\mu_1\mu_2\dots\mu_N}\} \quad (94)$$

by means of the relation [28, 29]

$$\gamma_M = e_M^{\mathbf{M}} \gamma_{\mathbf{M}} \quad (95)$$

in which we have introduced a vielbein of curved Clifford space C_{V_N} , given by the scalar product $\gamma_M^\dagger * \gamma^{\mathbf{M}}$. Notice a distinction between bold and normal indices, used for two different kinds of basis. The coordinate basis Clifford numbers $\gamma_M = \gamma_{\mu_1\mu_2\dots\mu_r}$ in general are *not* defined as a wedge product $\gamma_{\mu_1} \wedge \gamma_{\mu_2} \wedge \dots \wedge \gamma_{\mu_r}$. In particular, γ_M can be multivectors of definite grade, i.e., defined as a wedge product, but such property can hold only *locally* at a given point $\mathcal{E} \in C_{V_N}$, and cannot be preserved globally at all point $\mathcal{E}$ of our curved Clifford space. The relations for γ_M and the metric G_{MN} , discussed in sec. 4, refer to flat

C -space and are no longer generally valid in curved C -space. At a fixed point $\mathcal{E} \in C_{V_N}$ we can choose a coordinate system such that $\gamma_M = \gamma_{\mathbf{M}}$, and then the relations of Sec. 4 refer to flat C -space, spanned by $\gamma_{\mathbf{M}}$, i.e., the tangent space $T_{\mathcal{E}}(C_{V_N})$.

The set $\{\gamma_M\}$ of 2^n linearly independent coordinate basis fields (which depend on coordinates x^M) will be called a *coordinate frame field* in C -space.

The set $\{\gamma_{\mathbf{M}}\}$ of 2^n linearly independent orthonormal basis fields (which also in general depend on x^M) will be called *orthonormal frame field* in C -space.

Corresponding to each field γ_M we define a differential operator—which we call *derivative*— ∂_M , whose action depends on the quantity it acts on¹²:

(i) ∂_M maps scalars ϕ into scalars

$$\partial_M \phi = \frac{\partial \phi}{\partial x^M} \quad (96)$$

Then ∂_M is just the ordinary *partial derivative*.

(ii) ∂_M maps Clifford numbers into Clifford numbers. In particular, it maps a coordinate basis Clifford number γ_N into another Clifford number which can, of course, be expressed as a linear combination of γ_J :

$$\partial_M \gamma_N = \Gamma_{MN}^J \gamma_J \quad (97)$$

The above relation defines the connection Γ_{MN}^J for the coordinate frame field $\{\gamma_M\}$.

An analogous relation we have for the orthonormal frame field:

$$\partial_M \gamma_{\mathbf{M}} = -\Omega_{\mathbf{M}}^{\mathbf{N}} \gamma_{\mathbf{N}} \quad (98)$$

where $\Omega_{\mathbf{M}}^{\mathbf{N}}$ is the connection for the orthonormal frame field $\{\gamma_{\mathbf{M}}\}$.

When the derivative ∂_M acts on a polyvector valued field $A = A^N \gamma_N$ we obtain

$$\partial_M (A^N \gamma_N) = \partial_M A^N \gamma_N + A^N \partial_M \gamma_N = (\partial_M A^N + \Gamma_{MK}^N A^K) \gamma_N \equiv D_M A^N \gamma_N \quad (99)$$

where $D_M A^N \equiv \partial_M A^N + \Gamma_{MK}^N A^K$ are components of the covariant derivative in the coordinate basis, i.e., the ‘covariant derivative’ of the tensor analysis.

Here A^N are *scalar* components of A , and $\partial_M A^N$ is just the ordinary partial derivative with respect to x^M :

$$\partial_M \equiv \left(\frac{\partial}{\partial s}, \frac{\partial}{\partial x^{\mu_1}}, \frac{\partial}{\partial x^{\mu_1 \mu_2}}, \frac{\partial}{\partial x^{\mu_1 \dots \mu_n}} \right) \quad (100)$$

The derivative ∂_M behaves as a *partial derivative* when acting on scalars, and it defines a *connection* when acting on a basis $\{\gamma_M\}$. It has turned out very practical¹³ to use the

¹² This operator is a generalization to curved C -space of the derivative ∂_μ which acts in an n -dimensional space V_n , and was defined by Hestenes [9] (who used a different symbol, namely $\square_\mu$).

¹³ Especially when doing long calculation (which is usually the job of a theoretical physicist) it is much easier and quicker to write ∂_M than $\square_M$, ∇_M , D_{γ_M} , ∇_{γ_M} which all are symbols used in the literature.

easily writable symbol ∂_M which —when acting on a polyvector— cannot be confused with partial derivative. In ref. [29] we provided some arguments, why also conceptually it is more appropriate to retain the same symbol ∂_M , regardless of whether it acts on scalar, vector, or generic polyvector fields.

The derivative ∂_M is defined with respect to a coordinate frame field $\{\gamma_M\}$ in C -space. We can define a more fundamental derivative ∂ by

$$\partial = \gamma^M \partial_M \quad (101)$$

This is the *gradient* in C -space and it generalizes the ordinary gradient $\gamma^\mu \partial_\mu$, $\mu = 0, 1, 2, \dots, n-1$, discussed by Hestenes [9].

Besides the basis elements γ_M and $\gamma^{\mathbf{M}}$, we can define the reciprocal elements γ^M , $\gamma^{\mathbf{M}}$ by the relations

$$(\gamma^M)^\dagger * \gamma_N = \delta^M_N, \quad (\gamma^{\mathbf{M}})^\dagger * \gamma^{\mathbf{N}} = \delta^{\mathbf{M}}_{\mathbf{N}} \quad (102)$$

Curvature. We define the curvature of C -space in the analogous way as in the ordinary spacetime, namely by employing the commutator of the derivatives [9, 12, 35]. Using eq. (97) we have

$$[\partial_M, \partial_N] \gamma_J = R_{MNJ}{}^K \gamma_K \quad (103)$$

where

$$R_{MNJ}{}^K = \partial_M \Gamma_{NJ}^K - \partial_N \Gamma_{MJ}^K + \Gamma_{NJ}^R \Gamma_{MR}^K - \Gamma_{MJ}^R \Gamma_{NR}^K \quad (104)$$

is the curvature of C -space. Using (103) we can express the curvature according to

$$(\gamma^K)^\dagger * ([\partial_M, \partial_N] \gamma_J) = R_{MNJ}{}^K \quad (105)$$

An analogous relation we have if the commutator of the derivatives operates on a orthonormal basis elements and use eq. (98)

5.2 On the dynamical curved C -space with sources

We will assume that physics takes place in curved C -space. The latter space is a generalization of curved spacetime. As in the ordinary general relativity the metric $g_{\mu\nu}$ is a dynamical quantity, so in the generalized general relativity the C -space metric G_{MN} is a dynamical quantity. Instead of the usual point particles and branes the role of the sources is now assumed by the generalized branes (which include the generalized point particles) described by one of the classically equivalent actions (69),(73) or (76), which we will denote by a common symbol I_{Brane} .

The action thus contains a term which describes a C -space brane and a kinetic term which describes the dynamics of the C -space itself:

$$I = I_{\text{Brane}} + \frac{1}{16\pi G_C} \int d^{2n}x \sqrt{|G|} R \quad (106)$$

Here G_C is a constant (having the role of the “gravitational” constant in C -space), $G \equiv \det G_{MN}$ the determinant of the C -space metric, and $R = R_{MNJK}G^{MJ}G^{NK}$ the curvature scalar of C -space.

More details about how to proceed with and further generalize the theory based on the action (106) are to be found in [12]. The idea that the curved C -space can provide a realization of Kaluza-Klein theory has been considered in refs. [38, 41, 28, 29]. In this paper we will explore the brane part I_{Brane} which contains the coupling of the C -space brane’s embedding functions $X^M(\xi^A)$ to the C -space metric G_{MN} .

5.3 The generalized branes in curved C -space

We will consider a generalized brane (a C -space brane) moving in a fixed curved C -space background. We will assume that the action is given by eq. (73) in which there occurs the metric G_{MN} of the curved target C -space into which our generalized brane is embedded.

Suppose that in the target C -space there exist isometries described by K Clifford numbers $k^\alpha = k^\alpha_M \gamma^M$, $\alpha = 1, 2, \dots, K$; $M = 1, 2, \dots, 16$, (generalized Killing “vectors”), whose components satisfy the conditions

$$D_M k^\alpha_N + D_N k^\alpha_M = 0 \quad (107)$$

where the covariant derivative $D_M A_N$ of components A_M of an arbitrary polyvector A is given in eq. (99).

Splitting the coordinate basis and orthonormal basis indices according to

(i) *coordinate basis indices*: $M = (\mu, \bar{M})$, $\mu = 0, 1, 2, 3$; $\bar{M} = 4, 5, \dots, 16$

(ii) *orthonormal basis indices*: $\mathbf{M} = (\boldsymbol{\mu}, \bar{\mathbf{M}})$, $\boldsymbol{\mu} = 0, 1, 2, 3$; $\bar{\mathbf{M}} = 4, 5, \dots, 16$

the metric and vielbein can be written as

$$G_{MN} = \begin{pmatrix} G_{\mu\nu} & G_{\mu\bar{M}} \\ G_{\bar{M}\nu} & G_{\bar{M}\bar{N}} \end{pmatrix}, \quad e^{\mathbf{M}}_M = \begin{pmatrix} e^{\boldsymbol{\mu}}_\mu & e^{\boldsymbol{\mu}}_{\bar{M}} \\ e^{\bar{\mathbf{M}}}_\mu & e^{\bar{\mathbf{M}}}_{\bar{M}} \end{pmatrix} \quad (108)$$

Let us recall that that the vielbein according to eq. (84) can be written as the scalar product of the C -space coordinate and orthonormal basis elements:

$$e_M^{\mathbf{M}} = \gamma_M^\dagger * \gamma^{\mathbf{M}} \quad (109)$$

We can now assume that the orthonormal basis $\{\gamma_{\mathbf{M}}\}$ is chosen so that $\gamma_{\boldsymbol{\mu}}$, $\boldsymbol{\mu} = 0, 1, 2, 3$, are tangent vectors to the spacetime (a subspace of C -space), whilst the remaining basis elements $\gamma_{\bar{\mathbf{M}}}$, $\bar{\mathbf{M}} = 4, 5, \dots, 16$ are tangent to the “internal” part of C -space. Since the basis is orthogonal, we have

$$\gamma_{\boldsymbol{\mu}}^\dagger * \gamma_{\bar{\mathbf{M}}} = 0 \quad (110)$$

Next we assume that the coordinate basis $\{\gamma_M\}$ is chosen so that γ_μ , $\mu = 0, 1, 2, 3$ in general is *not* tangent to V_4 , whilst the remaining coordinate basis elements $\gamma_{\bar{M}}$, $\bar{M} =$

4, 5, ..., 16, are tangent to the “internal” part of C -space. This means that $\gamma_\mu * \gamma_{\bar{M}} \neq 0$, whilst

$$\gamma_\mu^\dagger * \gamma_{\bar{M}} = 0 \quad (111)$$

The latter equation can be written as $\eta_{\mu\nu} \gamma^{\nu\dagger} * \gamma_{\bar{M}} = \eta_{\mu\nu} e^\nu_{\bar{M}} = 0$. Since $\eta_{\mu\nu}$ is diagonal, it follows that

$$e^\mu_{\bar{M}} = 0 \quad (112)$$

Taking a coordinate system in which

$$k^{\alpha\mu} = 0, \quad k^{\alpha\bar{M}} \neq 0 \quad (113)$$

the components $e^{\bar{M}}_\mu$ can be written in terms of the Killing vectors and gauge fields $A_\mu^\alpha(x^\mu)$:

$$e^{\bar{M}}_\mu = e^{\bar{M}}_{\bar{M}} k^{\alpha\bar{M}} A_\mu^\alpha; \quad \partial_{\bar{M}} A_\mu^\alpha = 0 \quad (114)$$

For the “mixed” components of the inverse vielbein we find

$$e^\mu_{\bar{M}} = 0 \quad (115)$$

This follows directly from

$$\gamma^{\mu\dagger} * \gamma_{\bar{M}} = 0 = \gamma^{\mu\dagger} (e^{\bar{M}}_{\bar{M}} \gamma_{\bar{M}}) = \gamma^{\mu\dagger} * (e^\mu_{\bar{M}} \gamma_\mu + e^{\bar{M}}_{\bar{M}} \gamma_{\bar{M}}) = e^\mu_{\bar{M}} e^{\bar{M}}_{\bar{M}} \quad (116)$$

where we have used eq. (112). Since in general $e^{\bar{M}}_{\bar{M}} \neq 0$, it follows that $e^\mu_{\bar{M}}$ is equal to zero.

Using (95), (112) and (115) and the above choice of coordinates and local orthonormal frame we find that the quadratic form can be written as the sum of the 4-dimensional quadratic form and the part due to the “extra dimensions”:

$$dX^\dagger * dX = (dx^M \gamma_M)^\dagger * (dx^N \gamma_N) = g_{\mu\nu} dx^\mu dx^\nu + \phi^{\bar{M}\bar{N}} dx_{\bar{M}} dx_{\bar{N}} \quad (117)$$

where

$$g_{\mu\nu} = e^\mu_\mu e^\nu_\nu \gamma_\mu \gamma_\nu = G_{\mu\nu} - \phi^{\bar{M}\bar{N}} k^\alpha_{\bar{M}} k^\beta_{\bar{N}} A_\mu^\alpha A_\nu^\beta \quad \text{and} \quad \phi^{\bar{M}\bar{N}} \equiv e^{\bar{M}}_{\bar{M}} e^{\bar{N}}_{\bar{N}} \eta^{\bar{M}\bar{N}} \quad (118)$$

Here $G_{\bar{M}\bar{N}} \equiv \phi_{\bar{M}\bar{N}} = e^{\bar{M}}_{\bar{M}} e^{\bar{N}}_{\bar{N}} \eta_{\bar{M}\bar{N}}$, and $\phi^{\bar{M}\bar{N}}$ is the inverse of $\phi_{\bar{M}\bar{N}}$ in the “internal” space. Notice the validity of eqs. (112) and (115).

Let us now use the splitting (117) in the brane action (73). We obtain

$$\partial_A X^M \partial_B X^N G_{MN} = \partial_A X^\mu \partial_B X^\nu g_{\mu\nu} + \partial_A X_{\bar{M}} \partial_B X_{\bar{N}} \phi^{\bar{M}\bar{N}} \quad (119)$$

The auxiliary variables G_{AB} and the induced metric on the (generalized, i.e., C -space) brane are related according to eq. (75). Let us introduce new auxiliary variables G'_{AB} and new brane tension T' according to

$$G_{AB} = G'_{AB} + \partial_A X_{\bar{M}} \partial_B X_{\bar{N}} \Phi^{\bar{M}\bar{N}} \quad (120)$$

$$T\sqrt{|G|}G^{AB} = T'\sqrt{|G'|}G'^{AB} \quad (121)$$

where G^{AB} and G'^{AB} are the inverse matrices of G_{AB} and G'_{AB} , respectively.

Using eqs. (120),(121) we have

$$T\sqrt{|G|}d_C = T\sqrt{|G|}G^{AB}G_{AB} = T'\sqrt{|G'|}G'^{AB}(G'_{AB} + \partial_A X_{\bar{M}}\partial_B X_{\bar{N}}\Phi^{\bar{M}\bar{N}}) \quad (122)$$

$$T\sqrt{|G|} = \frac{T\sqrt{|G|}d_C}{d_C} = \frac{T\sqrt{|G|}G^{AB}G_{AB}}{d_C} = \frac{T'\sqrt{|G'|}}{d_C}G'^{AB}(G'_{AB} + \partial_A X_{\bar{M}}\partial_B X_{\bar{N}}\Phi^{\bar{M}\bar{N}}) \quad (123)$$

From eqs. (122),(123) we find that the extra term in the brane action (73) can be written

$$T\sqrt{|G|}(2 - d_C) = T'\sqrt{|G'|}(2 - d_C) + P^A_{\bar{M}}\partial_A X_{\bar{N}}\Phi^{\bar{M}\bar{N}}\left(\frac{2}{d_C} - 1\right) \quad (124)$$

where we have written

$$T\sqrt{|G|}G^{AB}\partial_A X_{\bar{M}} = T'\sqrt{|G'|}G'^{AB}\partial_A X_{\bar{M}} = P^A_{\bar{M}} \quad (125)$$

Here $P^A_{\bar{M}}$ are extra components of the brane canonical momentum $P^A_{\bar{M}} = \partial\mathcal{L}/\partial\partial_B X^M = T\sqrt{|G|}G^{AB}\partial_A X_M$.

Inserting eqs. (120),(121) and (124) into the brane action (73) we obtain

$$\begin{aligned} I_{\text{Brane}}[X^\mu, G'^{AB}] &= \frac{1}{2} \int d^{d_C} \xi T' \sqrt{|G'|} (G'^{AB} \partial_A X^\mu \partial_B X^\nu g_{\mu\nu} + 2 - d_C) \\ &\quad + \frac{1}{d_C} \int d^{d_C} \xi P^A_{\bar{M}} \partial_A X_{\bar{N}} \Phi^{\bar{M}\bar{N}} \end{aligned} \quad (126)$$

Variation of the latter action with respect to G'^{AB} gives

$$G'_{AB} = \partial_A X^\mu \partial_B X^\nu g_{\mu\nu} \quad (127)$$

which is consistent with eqs. (75),(119),(120). Since the index μ runs over the embedding spacetime coordinates $\mu = 0, 1, 2, \dots, N-1$ and $A, B = 1, 2, \dots, d_C$ run over the coordinates of the brane's C -space worldsheet C_{V_n} of dimension $d_C = 2^n$, we have that the system of equations (127) is determined, if $2^n \leq N$. In the case $N = 4$ we have that $2^n \leq 4$, which is satisfied for $n \leq 2$. In 4-dimensional spacetime the splitting described above works for point particles ($n = 1$) and strings ($n = 2$). Higher dimensional branes, that do not satisfy $2^n \leq N$, of course, can also exist, but G'_{AB} then cannot have the role of auxiliary variables, because the system of equations (127) is then underdetermined. In order to obtain a determined system for auxiliary variables, a splitting of the indices A, B , analogous to that of M, N is then needed as well.

In eq. (126) we have an action for a generalized C -space brane worldsheet C_{V_n} embedded in a higher dimensional curved manifold (which in our case is the embedding C -space

C_{V_N}). In particular, if we have just a usual point particle sweeping a 1-dimensional ‘world-sheet’, i.e., a worldline, embedded in a higher dimensional curved space, which can be either a C -space or just simply a spacetime with extra dimensions, then we have to take $d_C = 1$, and the action (126) becomes

$$I[X^\mu, \Lambda] = \frac{M}{2} \int d\tau \Lambda \left(\frac{\dot{X}^\mu \dot{X}^\nu g_{\mu\nu}}{\Lambda^2} + 1 \right) + \int d\tau P_{\bar{M}} \dot{X}_{\bar{N}} \Phi^{\bar{M}\bar{N}} \quad (128)$$

Above we have denoted

$$\begin{aligned} G'_{AB} &= G'_{11} \equiv \Lambda^2, & G'^{AB} &= G'^{11} \equiv \frac{1}{\Lambda^2}, & T' &\equiv M \\ \partial_A X^M &= \partial_1 X^M \equiv \frac{dX^M}{d\tau} \equiv \dot{X}^M \\ P^A_{\bar{M}} &= P^1_{\bar{M}} \equiv P_{\bar{M}} \end{aligned} \quad (129)$$

The action (128) can also be obtained by splitting the following point particle action:

$$I[X^M, \lambda] = \frac{m}{2} \int d\tau \lambda \left(\frac{\dot{X}^M \dot{X}^N G_{MN}}{\lambda^2} + 1 \right) \quad (130)$$

Variation of the latter action with respect to λ gives

$$\lambda^2 = \dot{X}^M \dot{X}_M \quad (131)$$

which can be split according to

$$\dot{X}^M \dot{X}_M = \dot{X}^\mu \dot{X}^\nu g_{\mu\nu} + \dot{X}_{\bar{M}} \dot{X}_{\bar{N}} \Phi^{\bar{M}\bar{N}} \quad (132)$$

Introducing a new auxiliary variable Λ and new mass M (a 4-dimensional mass) according to

$$\lambda^2 = \Lambda^2 + \dot{X}_{\bar{M}} \dot{X}_{\bar{N}} \Phi^{\bar{M}\bar{N}} \quad (133)$$

$$\frac{m}{\lambda} = \frac{M}{\Lambda} \quad (134)$$

and inserting eqs. (132)–(134) into the point particle action (130) we obtain the split point particle action (128), where

$$P_{\bar{M}} = \frac{m}{\lambda} \dot{X}_{\bar{M}} = \frac{M}{\Lambda} \dot{X}_{\bar{M}} \quad (135)$$

So we have verified that the split generalized brane action (126) includes a correct point particle action. Relations (132)–(135) above are point particle analog of the brane relations (119)–(121), (125).

Let us now return to the generalized brane action (126). The extra term can be written as

$$\begin{aligned} P^A_{\bar{M}} \partial_A X_{\bar{N}} \Phi^{\bar{M}\bar{N}} &= P^A_{\bar{M}} \Phi^{\bar{M}\bar{N}} G_{\bar{N}J} \partial_A X^J = P^A_{\bar{M}} \Phi^{\bar{M}\bar{N}} A^\alpha_J k_{\bar{N}\alpha} \partial_A X^J \\ &= J^A_\alpha A^\alpha_J \partial_A X^J = J^A_\alpha (A^\alpha_\mu \partial_A X^\mu + A^\alpha_{\bar{M}} \partial_A X^{\bar{M}}) \end{aligned} \quad (136)$$

where

$$J^A_\alpha \equiv P^A_{\bar{M}} \Phi^{\bar{M}\bar{N}} k_{\bar{N}\alpha} = P^A_{\bar{M}} k^{\bar{M}}_\alpha = P^A_M k^M_\alpha \quad (137)$$

are current densities which are conserved due to the presence of isometries. The last step in eq. (137) holds because of eq. (113). In eq. (136) we have written the metric components in terms of gauge potentials and Killing “vectors”

$$G_{\bar{N}J} = A^\alpha_J k_{\bar{N}\alpha} \quad (138)$$

In particular,

$$G_{\bar{N}\mu} = A^\alpha_\mu k_{\bar{N}\alpha} \quad \text{if } J = \mu \quad (139)$$

$$G_{\bar{N}\bar{M}} = A^\alpha_{\bar{M}} k_{\bar{N}\alpha} \quad \text{if } J = \bar{M} \quad (140)$$

In eq. (139) we have the well known relation between gauge potentials, Killing “vectors” and “mixed” components of the metric, the relation that was derived (see e.g., [39, 40]) within the context of an ordinary higher dimensional spacetime manifold. In eq. (140) we have rewritten the “internal” space metric components in terms of gauge potentials $A^\alpha_{\bar{M}}$ and Killing “vectors”.

Inserting eq. (136) into eq. (126) we obtain

$$\begin{aligned} I_{\text{Brane}}[X^\mu, G'^{AB}] &= \frac{1}{2} \int d^d \xi T' \sqrt{|G'|} (G'^{AB} \partial_A X^\mu \partial_B X^\nu g_{\mu\nu} + 2 - d_C) \\ &\quad + \frac{1}{d_C} \int d^d \xi J^A_\alpha A^\alpha_M \partial_A X^M \end{aligned} \quad (141)$$

The last term is just the interactive term between the conserved current densities J^A_α and gauge potentials A^α_M coupled to

$$\partial_A X^M \equiv \partial_{a_1 \dots a_r} X^{\mu_1 \dots \mu_R}, \quad r = 0, 1, 2, \dots, n; \quad R = 0, 1, 2, \dots, N \quad (142)$$

The latter potentials include the ordinary nonabelian potentials A^α_μ , $\alpha = 1, 2, \dots, K$, coupled to $\partial_A X^\mu$, and the extra potentials $A^\alpha_{\bar{M}}$, coupled to $\partial_A X^{\bar{M}}$. In sec. we argued that in a particular parametrization of $\xi^A = (\xi, \xi^a, \xi^{a_1 a_2}, \dots)$, if we take $r = R = 0, 1, 2, \dots, n$, we can set

$$\partial_{a_1 \dots a_r} X^{\mu_1 \dots \mu_r} = \partial_{[a_1} X^{\mu_1} \dots \partial_{a_r]} X^{\mu_r} \quad (143)$$

If $r = 1$, the above relation is automatically true, because we have just $\partial_{a_1} X^{\mu_1}$ on both sides of the equation. If $r = 0$, then we have derivative with respect to scalar parameter

ξ , and eq. (143) is automatically satisfied as well. But for higher grades, $r = 2, 3, \dots, n$, eq. (143) must be imposed as an extra condition, namely a condition that fixes a gauge.

If we assume the validity of relation (143), we find that the interactive term in eq. (141), namely

$$\begin{aligned} I_{\text{int}} &= \frac{1}{d_C} \int d^{d_C} \xi J^A_{\alpha} A^{\alpha}_M \partial_A X^M = \frac{1}{d_C} \int d^{d_C} \xi J^A_{\alpha} (A^{\alpha}_{\mu} \partial_A X^{\mu} + A^{\alpha}_{\bar{M}} \partial_A X^{\bar{M}}) \\ &= \frac{1}{d_C} \int d^{d_C} \xi J^{\alpha_1 \dots \alpha_r}_{\alpha} A^{\alpha}_{\mu_1 \dots \mu_R} \partial_{a_1 \dots a_r} X^{\mu_1 \dots \mu_R} \end{aligned} \quad (144)$$

contains the coupling of the antisymmetric gauge potentials to the antisymmetric current density. In the case of a single Killing “vector” field, $\alpha = 1$, the gauge fields $A_{\mu_1 \dots \mu_R}$ are abelian, and for $R = r = n$, eq. (144) becomes the interactive term for the well known *Kalb-Ramond fields* [42]. The latter fields have an important role in string theories and brane theories [4]. Here we have demonstrated a possible broader theoretical framework for generalized branes, coupled to generalized gauge fields, which includes strings and Kalb-Ramond fields. An alternative approach to generalized gauge (Maxwell) fields in C -space has been considered in refs. [43].

To sum up, the action (141) contains the coupling of generalized gauge fields with the charge current density. Besides the ordinary gauge fields A^{α}_{μ} there also occur higher grade, Kalb-Ramond fields $A^{\alpha}_{\bar{M}} \equiv A^{\alpha}_{\mu_2 \dots \mu_R}$ and the zero grade, scalar, field A^{α} . All those fields are included in the compact coupling term (144), where $R = 0$ stands for the scalar component, $R = 1$ for vector and $R = 2, 3, \dots, N$ for higher grade components, and analogously for $r = 0, 1, 2, \dots, n$. Eq. (144) contains the non abelian generalization of the well known coupling term for Kalb-Ramond fields considered in refs. [42, 44, 8]. The latter coupling occurs as a special case of eq. (144), if we take the terms with the grade $r = R = n$ only, i.e., the terms with the maximal grade of the worldsheet multivectors. However, more general coupling terms with $r \neq R \neq n$ also exists in our theory. Thus the case for $r = n$, $R = 1$ includes the well known electrically charged closed membrane considered by Dirac [45]. In the following subsection we will discuss more explicitly some particular cases of our general theory, that were previously considered in the literature as separate subjects.

5.4 Comparison with previous theories of charged branes

The electromagnetic potential A_{μ} couples to the time like tangent element of the worldline $X^{\mu}(\tau)$ of a charged *particle*:

$$I_{\text{int}}^{\text{particle}} = q \int d\tau \frac{\partial X^{\mu}}{\partial \tau} A_{\mu} \quad (145)$$

where q is the particle’s electric charge.

For extended objects the coupling involves the worldsheet tangent elements (velocities) (27) and generalized Maxwell potentials $A_{\mu_1 \dots \mu_n}$. The corresponding equation for a *string* is

$$I_{\text{int}}^{\text{string}} = \frac{q^{12}}{2!} \int d^2 \xi \partial_{[1} X^\mu \partial_{2]} X^\nu A_{\mu\nu} \quad (146)$$

where the two-vector charge q^{12} is coupled to a two-vector potential $A_{\mu\nu}$. For a generic *brane* the coupling reads

$$I_{\text{int}}^{\text{brane}} = \frac{1}{n!} q^{12 \dots n} \int d^n \xi \partial_{[1} X^{\mu_1} \dots \partial_{n]} X^{\mu_n} A_{\mu_1 \dots \mu_n} \quad (147)$$

Eq.(147) can be rewritten as

$$I_{\text{int}}^{\text{brane}} = \left(\frac{1}{n!} \right)^2 q^{a_1 a_2 \dots a_n} \int d^n \xi \partial_{[a_1} X^{\mu_1} \dots \partial_{a_n]} X^{\mu_n} A_{\mu_1 \dots \mu_n} \quad (148)$$

Here $q, q^{a_1 a_2}, \dots, q^{a_1 a_2 \dots a_n}$ are the coupling strengths.

If we introduce

$$q_n \equiv \frac{1}{n!} q^{a_1 \dots a_n} \epsilon_{a_1 \dots a_n} \quad (149)$$

and use eqs. (46),(50) we have

$$I_{\text{int}}^{\text{brane}} = \frac{1}{n!} q_n \int d\tilde{\xi} \frac{\partial X^{\mu_1 \dots \mu_n}}{\partial \tilde{\xi}} A_{\mu_1 \dots \mu_n} \quad (150)$$

In the case considered above the vector potential A_μ couples to the vector tangent element of the particle's world line, the 2-vector potential $A_{\mu\nu}$ couples to the 2-vector tangent element of the string's world sheet, and in general, the n -vector potential $A_{\mu_1 \dots \mu_n}$ couples to the n -vector tangent element of the brane's worldsheet. By this the possible couplings are not exhausted. Long time ago Dirac considered a relativistic charged closed membrane coupled to the ordinary Maxwell field given in terms of the vector potential A_μ . Later, this theory has been generalized [46, 47] to closed branes of any dimension.

It is well known that for *open strings and membranes* the electric charge q —dues to its repulsive character— can only be concentrated on the boundary (e.g., at the string's ends). But for *closed strings* and, in general, closed p -branes, q can be distributed over such extended objects. So instead of the total charge q we have to introduce the *charge density*. But since the brane moves, we have the charge current density on the brane. In a covariant description we introduce the *charge current density* j^a on the brane's worldsheet V_n . In ref. [47] the following action was considered:

$$I[X^\mu, A_\mu] = \int d^n \xi (\kappa \sqrt{|\det \partial_a X^\mu \partial_b X_\mu|} + j^a \partial_a X^\mu A_\mu) + \frac{1}{4\pi} \int d^N x \sqrt{|g|} F_{\mu\nu} F^{\mu\nu} \quad (151)$$

The first term in eq.(151) is just the minimal surface Dirac-Nambu-Goto action for a p -brane ($n = p + 1$), whilst the second term represents the minimal coupling of the brane's

electric charge current density j^a with the electromagnetic field potential A_μ . The last term is the kinetic term for the electromagnetic field $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$.

The action (151) has the following transformation property:

$$I[X^\mu, A'_\mu] = I[X^\mu, A_\mu] + \int d^n \xi j^a \partial_a X^\mu \partial_\mu \varphi = I[X^\mu, A_\mu] + \int d^n \xi \partial_a (j^a \varphi) \quad (152)$$

where

$$A'_\mu = A_\mu + \partial_\mu \varphi \quad (153)$$

and

$$\partial_a j^a = 0 \quad (154)$$

Eq.(4.8) is the gauge transformation of A_μ , whilst eq.(154) expresses the charge conservation. So we have

$$\int d^n \xi \partial_a j^a = \oint d\Sigma_a j^a = q(\Sigma_2) - q(\Sigma_1) = 0 \quad (155)$$

where $q = \int d\Sigma_a j^a$ is the total charge of the closed brane and $d\Sigma_a$ the hypersurface element. Here j^a is a worldsheet vector density (not vector), so that (155) is covariant under reparametrizations of V_n .

Here j^a is the *intrinsic current density*, a vector density on V_n . By the relation

$$j^\mu = \int d^n \xi \delta(x - X(\xi)) j^a \partial_a X^\mu \quad (156)$$

we obtain the *extrinsic current density*, i.e., the current density in the target space V_N . The minimal coupling Lagrangian in (151) then reads

$$\int d^n \xi j^a \partial_a X^\mu A_\mu = \int d^N x j^\mu A_\mu \quad (157)$$

It is straightforward to show that the conservation law (154) implies

$$\partial_\mu j^\mu = 0 \quad (158)$$

The transformed action (which is a functional of X^μ and A'_μ) differs from the “original” action $I[X^\mu, A_\mu]$ by a term with the total divergence which has no influence on the equations of motion. The *canonical momentum density* is

$$p_\mu{}^a = \frac{\mathcal{L}}{\partial \partial_a X^\mu} = \kappa \sqrt{|f|} \partial^a X_\mu + j^a A_\mu \quad (159)$$

and it obviously is not invariant under gauge transformations (153). Therefore it is convenient to introduce the *kinetic momentum density*

$$\pi_\mu{}^a \equiv p_\mu{}^a - j^a A_\mu \quad (160)$$

which is gauge invariant. By employing (160) we can write *the phase space action* [46]

$$I_{\text{Brane}}[X^\mu, p_\mu^a, \gamma_{ab}] = \int d^n \xi \left[p_\mu^a \partial_a X^\mu - \frac{1}{2} \frac{\gamma_{ab}}{\kappa \sqrt{|\gamma|}} (\pi_\mu^a \pi_\nu^b g^{\mu\nu} - \kappa^2 |\gamma| \gamma^{ab}) + \kappa \sqrt{|\gamma|} (1 - n) \right] \quad (161)$$

After eliminating p_μ^a from its equations of motion

$$p_c^\nu - j_c A^\nu = \kappa \sqrt{|\gamma|} \partial_c X^\nu \quad (162)$$

we obtain [47, 12] the Howe-Tucker action in the presence of electromagnetic field:

$$I_{\text{Brane}}[X^\mu, \gamma_{ab}] = \int d^n \xi \left[\frac{\kappa \sqrt{|\gamma|}}{2} (\gamma^{ab} \partial_a X^\mu \partial_b X_\mu + 2 - n) + j^a \partial_a X^\mu A_\mu \right] \quad (163)$$

The total action is thus the sum of the brane action I_{Brane} and the kinetic term for the electromagnetic field I_{EM} (given by the last term in (151)).

A closed brane can thus possess two different kinds of charges coupled to two different kinds of gauge fields:

- (i) Kalb-Ramond charge $q^{12\dots n}$ coupled to Kalb-Ramond gauge field potentials $A_{\mu_1\dots\mu_n}$ according to eq.(147);
- (ii) the ordinary electric charge q coupled to the ordinary Maxwell potential A_μ according to eq.(151) (or equivalently, eqs. (161) and (163)).

By inspecting the interactive action (147) we observe that it is just a coupling term that can be added to the action (40). On the other hand, the action (163) is just a generalization of the action (28) to which we added the minimal coupling term for the Maxwell field A_μ . Both actions, (28) and (40), were generalized to C -space, and so we have obtained the action (69) (which is equivalent to the action (73)). As we have seen in Sec. 5.3, the two kinds of couplings, namely (i) and (ii), can be unified by employing the C -space description, in which they arise from the metric of the target C -space C_{V_N} , equipped with connection, whose curvature was assumed to be in general different from zero.

6 Discussion

Spacetime as a continuum of points is just a start, from which we can arrive at a continuum of oriented areas, also called oriented volumes. Such enlarged continuum, called Clifford space or C -space provides a framework for generalizing the concepts of event, point particle, string and, in general, brane. As an ordinary brane is an object that extends in

spacetime, so a generalized brane is an object that extends in C -space. An important feature of C -space is that oriented areas (volumes) of different grades, associated with branes of different dimensionalities, can be transformed into each other. A generic geometric object has mixed grade, it is represented by a Clifford number, also called Clifford aggregate or polyvector, and it is associated with a generalized brane. Having set such a kinematics, one can construct a dynamics which employs the well known Howe-Tucker brane action, which is now generalized to C -space. The latter space can be curved, and this gives rise, à la Kaluza-Klein, to gauge fields as components of the C -space metric. The C -space Howe-Tucker action, minimally coupled to the C -space metric, contains, in particular, the well known terms for the coupling of a p -brane (including a point particle) to gauge fields. Amongst the latter gauge fields there also occur the Kalb-Ramond anti-symmetric fields, and their non Abelian generalizations [48]. In this paper we focused our attention to the dynamics of the branes, and left aside the fact that in a more complete treatment [28, 29] C -space itself becomes dynamical, as in general relativity.

In the present paper we presented only a piece of the story, namely, how the classical theory of branes in a fixed background could be generalized to Clifford space. Since with the points of a flat Clifford space one can associate Clifford numbers (polyvectors), that are elements of Clifford algebra $\mathbf{C}_N$, this automatically brings spinors (as members of left or right ideals of $\mathbf{C}_N$) into our description. A polyvector $X^{\mu_1 \dots \mu_R} \gamma_{\mu_1 \dots \mu_R}$, since it can be rewritten, e.g., in terms of a basis spanning all independent left ideals, thus contains spinorial degrees of freedom [28, 29, 16]. This means that by describing our branes in terms of the C -space embedding functions $X^M \equiv X^{\mu_1 \dots \mu_R}$ we have already included spinorial degrees of freedom. We do not need to postulate them separately, as is done in ordinary string and brane theories, where besides Grassmann even (‘‘bosonic’’) variables X^μ , there occur also Grassmann odd (‘‘fermionic’’) or spinorial variables. In this formulation we have a possible clue to the resolution of a big open problem, namely, what exactly is string theory. I believe that further research into that direction would provide very fruitful results and insight. An important insight is already in recognizing that 16-dimensional Clifford space provides a consistent description of quantized string theory [38, 16]. The underlying spacetime can remain 4-dimensional, there is no need for a 26-dimensional or a 10-dimensional spacetime. The extra degrees of freedom required for consistency of string theory, described in terms of variables $X^M(\tau, \sigma)$, are due to extra dimensions of C -space, and they need not be compactified; they are due to volume (area) evolution, and are thus all physical. But since a generic component $X^M(\tau, \sigma) \equiv X^{\mu_1 \dots \mu_R}(\tau, \sigma)$ denotes an oriented R -volume, associated with an $(R - 1)$ -brane (i.e., a p -brane for $p = R - 1$), we have that string itself (i.e., 1-brane) is not enough for consistency. Higher branes are automatically present in the description with functions $X^{\mu_1 \dots \mu_R}(\tau, \sigma)$, although they are not described in full detail, but only up to the knowledge of oriented R -volume. Because

of the presence of two parameters τ , σ , we keep on talking about evolving *strings*, not in 26 or 10-dimensional spacetime, but in 16-dimensional Clifford space. In general, the number of parameters can be arbitrary, but less than 16, and so we have a brane in C -space, i.e., a generalized brane discussed in this paper.

The C -space approach to branes is possibly related to the approach with extra coordinates, considered by Duff and Lu [5] in order to describe brane dualities. Such interesting relation needs to be investigated in future research.

Clifford algebras in infinite dimensions and in continuous dimensions are still an uncharted territory worth exploring. A pioneering step into that direction has been done in ref. [12], where a theory of generalized branes was formulated in terms of the generators of the infinite dimensional Clifford algebra.

Another possible direction that remains to be further explored is related to the fact that p-brane actions can be recast as non Abelian gauge theories of volume preserving diffeomorphisms [8, 49]. It would be interesting to extend such approach to the case of the generalized branes in C -space considered in this paper. Also there is a number of works on polyvector generalized supersymmetries [50], on the implications to M theory [51], on generalized Yang-Mills theories, Poly-particles and duality in C -spaces [52, 48] and on quantum mechanics in C -spaces and non-commutativity [53]. A possible connection of C -space to twistors (see, e.g., [54]) would also be worth investigating.

Clifford algebra, without recourse to C -space and generalized branes, has been considered in numerous works attempting to explain the standard model [55]. There are diverse approaches, but common to all of them is a feeling that Clifford algebra might be a clue to the unification of fundamental forces. It is reasonable to expect that further research will bring useful results, and that it will be crucial to take into account the concepts of C -space and branes.

Acknowledgement

This work was supported by the Ministry of High Education, Science and Technology of Slovenia.

References

- [1] See, e.g., M.B. Green, J.H. Schwarz and E. Witten, *Superstring theory* (Cambridge University Press, Cambridge, 1987); M. Kaku, *Introduction to Superstrings* (Springer-Verlag, New York, 1988); U. Danielsson, *Rep. Progr. Phys.* **64**, 51 (2001).
- [2] J. Polchinski, "Lectures on D-branes," [arXiv:hep-th/9611050]; W. I. Taylor, "Lectures on D-branes, gauge theory and M(atrices)," arXiv:hep-th/9801182. H. Nicolai and R. Helling, "Supermembranes and M(atric) theory," arXiv:hep-th/9809103; J.H.Schwarz, *Phys. Rep.* **315**, 107 (1999).

- [3] L. Randall and R. Sundrum, *Phys. Rev. Lett.* **83**, 4690 (1999) [arXiv:hep-th/9906064].
M. J. Duff, “Benchmarks on the brane,” arXiv:hep-th/0407175.
- [4] M. J. Duff, R. R. Khuri and J. X. Lu, *Phys. Rept.* **259**, 213 (1995) [arXiv:hep-th/9412184].
M. J. Duff and J. X. Lu, *Class. Quant. Grav.* **9**, 1 (1992). M. J. Duff and J. X. Lu, *Phys. Rev. Lett.* **66**, 1402 (1991).
- [5] M. J. Duff, *Nucl. Phys. B* **335**, 610 (1990). M. J. Duff and J. X. Lu, *Nucl. Phys. B* **347**, 394 (1990).
- [6] A. Schild, *Phys. Rev.* **D16**, 1722 (1977).
- [7] T. Eguchi, *Phys. Rev. Lett.* **44**, 126 (1980).
- [8] A. Aurilia, A. Smailagic and E. Spallucci, *Phys. Rev. D* **47**, 2536 (1993) [arXiv:hep-th/9301019]; A. Aurilia and E. Spallucci, *Class. Quant. Grav.* **10**, 1217 (1993) [arXiv:hep-th/9305020]; S. Ansoldi, A. Aurilia and E. Spallucci, *Phys. Rev. D* **53**, 870 (1996) [arXiv:hep-th/9510133].
- [9] D. Hestenes, *Space-Time Algebra* (Gordon and Breach, New York, 1966); D. Hestenes and G. Sobczyk, *Clifford Algebra to Geometric Calculus* (D. Reidel, Dordrecht, 1984).
- [10] P. Lounesto, *Clifford Algebras and Spinors* (Cambridge University Press, Cambridge, 2001); B. Jancewicz, *Multivectors and Clifford Algebra in Electrodynamics* (World Scientific, Singapore, 1988); R. Porteous, *Clifford Algebras and the Classical Groups* (Cambridge University Press, 1995); W. Baylis, *Electrodynamics, A Modern Geometric Approach* (Boston, Birkhauser, 1999); A. Lasenby and C. Doran, *Geometric Algebra for Physicists* (Cambridge U. Press, Cambridge 2002); *Clifford Algebras and their applications in Mathematical Physics, Vol 1: Algebras and Physics*, eds by R. Ablamowicz, B. Fauser; Vol 2: Clifford analysis. eds by J. Ryan, W. Sprosig (Birkhauser, Boston, 2000); A.M.Moya, V.V Fernandez and W.A. Rodrigues, *Int.J.Theor.Phys.* **40**, 2347 (2001) [arXiv: math-ph/0302007]; “Multivector Functions of a Multivector Variable” [arXiv: math.GM/0212223]; Multivector Functionals [arXiv: math.GM/0212224]; W.A. Rodrigues, Jr, J. Vaz, Jr, *Adv. Appl. Clifford Algebras* **7**, 457 (1997); E.C de Oliveira and W.A. Rodrigues, Jr, *Ann. der Physik* **7**, 654 (1998). *Phys. Lett A* **291**, 367 (2001). W.A. Rodrigues, Jr, J.Y.Lu, *Foundations of Physics* **27**, 435 (1997); S. Vacaru, P. Stavrinos, E. Gaburov and D. Gonta, “Clifford and Riemann-Finsler structures in geometric mechanics and gravity,” [arXiv:gr-qc/0508023].
- [11] M. Pavšič, *Found. Phys.* **33**, 1277 (2003) [arXiv:gr-qc/0211085].
- [12] M.Pavšič, *The Landscape of Theoretical Physics: A Global View; From Point Particle to the Brane World and Beyond, in Search of Unifying Principle* (Kluwer Academic, Dordrecht 2001).
- [13] S. Ansoldi, A. Aurilia, C. Castro and E. Spallucci, *Phys. Rev.* **D64**, 026003 (2001) [arXiv:hep-th/0105027].
- [14] A. Aurilia, S. Ansoldi and E. Spallucci, *Class. Quant. Grav.* **19**, 3207 (2002) [arXiv:hep-th/0205028].
- [15] C. Castro and M. Pavšič, *Progr. Phys.* **1**, 31 (2005).

- [16] M. Pavšič, *Found. Phys.* **35**, 1617 (2005) [arXiv:hep-th/0501222].
- [17] E.C.G. Stueckelberg, *Helv. Phys. Acta*, **14**, 322 (1941); **14**, 588 (1941); **15**, 23 (1942)
- [18] R. P. Feynman *Phys. Rev.* **84**, 108 (1951)
- [19] J. Schwinger, *Phys. Rev.* **82**, 664 (1951)
- [20] W. C. Davidon, *Physical Review* **97**,1131 (1955); **97**,1139 (1955)
- [21] L. P. Horwitz and C. Piron, *Helv. Phys. Acta*, **46**, 316 (1973); L. P. Horwitz and F. Rohrlich, *Physical Review D* **24**, 1528 (1981); **26**, 3452 (1982); L. P. Horwitz, R. I. Arshansky and A. C. Elitzur *Found. Phys* **18**, 1159 (1988); R. Arshansky, L. P. Horwitz and Y. Lavie, *Foundations of Physics* **13**, 1167 (1983); L. P. Horwitz, in *Old and New Questions in Physics, Cosmology, Philosophy and Theoretical Biology* (Editor Alwyn van der Merwe, Plenum, New York, 1983); L. P. Horwitz and Y. Lavie, *Physical Review D* **26**, 819 (1982); L. Burakovsky, L. P. Horwitz and W. C. Schieve, *Physical Review D* **54**, 4029 (1996); L. P. Horwitz and W. C. Schieve, *Annals of Physics* **137**, 306 (1981)
- [22] J.R.Fanchi, *Phys. Rev. D* **20**, 3108 (1979); see also the review J.R.Fanchi, *Found. Phys.* **23**, 287 (1993), and many references therein; J. R. Fanchi *Parametrized Relativistic Quantum Theory* (Kluwer, Dordrecht, 1993)
- [23] H.Enatsu, *Progr. Theor. Phys* **30**, 236 (1963); *Nuovo Cimento A* **95**, 269 (1986); F. Reuse, *Foundations of Physics* **9**, 865 (1979); A. Kyprianidis *Physics Reports* **155**, 1 (1987); R. Kubo, *Nuovo Cimento A* , 293 (1985); M. B. Mensky and H. von Borzeszkowski, *Physics Letters A* **208**, 269 (1995); J. P. Aparicio, F. H. Gaioli and E. T. Garcia-Alvarez, *Physical Review A* **51**, 96 (1995); *Physics Letters A* **200**, 233 (1995); L. Hannibal, *International Journal of Theoretical Physics* **30**, 1445 (1991); F. H. Gaioli and E. T. Garcia-Alvarez, *General Relativity and Gravitation* **26**,1267 (1994)
- [24] M. Pavšič, *Found. Phys.* **21**, 1005 (1991); M. Pavšič, *Nuovo Cim.* **A104**, 1337 (1991); *Doga, Turkish Journ. Phys.* **17**, 768 (1993)
- [25] M. Pavšič, *Found. Phys.* **26**, 159 (1996) [arXiv:gr-qc/9506057].
- [26] M. Pavšič, *Found. Phys.* **31**, 1185 (2001) [arXiv:hep-th/0011216];
- [27] M. Riesz, “Sur certaines notions fondamentales en théorie quantiques relativiste’, in *Dixième Congrès Math. des Pays Scandinaves, Copenhagen, 1946* (Jul. Gjellerups Forlag, Copenhagen, 1947), pp. 123-148; M. Riesz, *Clifford Numbers and Spinors*, E. Bolinder and P. Lounesto (eds.), (Kluwer 1993); S. Teitler, *Supplemento al Nuovo Cimento* **III**, 1 (1965) and references therein; *Supplemento al Nuovo Cimento* **III**, 15 (1965); *J. Math. Phys.* **7**, 1730 (1966); **7**, 1739 (1966); W.A. Rodrigues, Jr., *J. Math. Phys.* **45**, 2908 (2004).
- [28] M. Pavšič, *Phys. Lett.* **B614**, 85 (2005) [arXiv:hep-th/0412255].
- [29] M. Pavšič, “Spin gauge theory of gravity in Clifford space: A realization of Kaluza-Klein theory in 4-dimensional spacetime,” *Int. J. Mod. Phys. A* **21**, 5905 (2006). arXiv:gr-qc/0507053.
- [30] L.P. Eisenhart, *Riemannian Geometry* (Princeton University Press, Princeton, 1926).

- [31] H. Rund, *Invariant Theory of Variational Problems on Subspaces of a Riemannian Manifold* (Van Den Hoeck & Rupert, Göttingen, 1971).
- [32] C.W. Misner, K.S. Thorne and J.A. Wheeler, *Gravitation* (W.H. Freeman and Company, San Francisco, 1973).
- [33] M. Pavšič, *Class. Quant. Grav.* **20**, 2697 (2003) [arXiv:gr-qc/0111092].
- [34] P.S. Howe and R.W. Tucker, *J. Phys. A: Math. Gen.* **10**, L155 (1977); A. Sugamoto, *Nucl. Phys. B* **215**, 381 (1983); M. Pavšič, *Class. Quant. Grav.* **5**, 247 (1988).
- [35] C. Castro and M. Pavšič, *Phys. Lett.* **B539**, 133 (2002) [arXiv:hep-th/0110079].
- [36] W. Pezzaglia, “Physical Applications of a Generalized Geometric Calculus,” in *Dirac Operators in Analysis* (Pitman Research Notes in Mathematics, Number 394), J. Ryan and D. Struppa eds., (Longmann 1997) pp.191–202 [arXiv: gr-qc/9710027]; “Dimensionally Democratic calculus and Principles of Polydimensional Physics,” in *Clifford Algebras and their Applications in Mathematical Physics*, R. Ablamowicz and B. Fauser eds. (Birkhauser 2000), pp.101–123,[arXiv: gr-qc/9912025]; “Classification of Multivector Theories and Modifications of the Postulates of Physics”, in *Clifford Algebras and their Applications in Mathematical Physics*, F. Brackx, R. Delanghe & H. Serras eds., (Kluwer 1993) pp.317–323, [arXiv: gr-qc/9306006].
- [37] C. Castro, *Chaos, Solitons and Fractals* **10**, 295 (1999); **11**, 1663 (2000); **12**, 1585 (2001); “The Search for the Origins of M Theory: Loop Quantum Mechanics, Loops/Strings and Bulk/Boundary Dualities”, arXiv: hep-th/9809102; C. Castro, *Found. Phys.* **30**, 1301 (2000).
- [38] M. Pavšič, “Clifford space as a generalization of spacetime: Prospects for unification in physics,” arXiv:hep-th/0411053.
- [39] J.F Luciani, *Nucl. Phys.* **B135**, 111 (1978).
- [40] E. Witten, *Nucl. Phys.* **B186**, 412 (1981).
- [41] C. Castro, *Found. Phys.* **35**, 971 (2005).
- [42] M. Kalb and P. Ramond, *Phys. Rev.* **D9**, 2273 (1974).
- [43] C. Castro, *Mod. Phys. Lett. A* **19**, 19 (2004).
- [44] A. Aurilia and Y. Takahashi, *Progr. Theor. Phys.* **66** 693 (1981).
- [45] P.A.M.Dirac, *Proc.R.Soc.(London) A* **268**, 57 (1962).
- [46] A. O. Barut and M. Pavšič, *Mod. Phys. Lett. A* **7**, 1381 (1992).
- [47] A. O. Barut and M. Pavšič, *Phys. Lett. B* **306**, 49 (1993); *Phys. Lett. B* **331**, 45 (1994).
- [48] G. Savidy “Non-Abelian tensor gauge fields: Enhanced symmetries,” arXiv:hep-th/0604118.
- [49] C. Castro, *Int. J. Mod. Phys. A* **13**, 1263 (1998)

- [50] C. Castro, *Int. J. Mod. Phys. A* **21**, 2149 (2006). D. V. Alekseevsky, V. Cortes, C. Devchand and A. Van Proeyen, *Commun. Math. Phys.* **253**, 385 (2004) [arXiv:hep-th/0311107].
- [51] C. Castro, *J. Math. Phys.* **47**, 112301 (2006); Z. Kuznetsova and F. Toppan, arXiv:hep-th/0610122.
- [52] C. Castro, *Annals Phys.* **321**, 813 (2006). C. Castro, *Found. Phys.* **34**, 1091 (2004). M. Land, *Found. Phys.* **35**, 1245 (2005) [arXiv:hep-th/0603169].
- [53] C. Castro, *J. Phys. A: Math. Gen.* **39**, 14205 (2006)
- [54] R. Penrose, *Chaos, Solitons & Fractals* **10**, 581 (1999)
- [55] F. Smith, *Intern. J. Theor. Phys.* **24**, 155 (1985); **25**, 355 (1985); G. Trayling and W.E. Baylis, *Int. J. Mod. Phys. A* **16**, Suppl. 1C (2001) 900; *J. Phys. A: Math. Gen.* **34**, 3309 (2001); G. Roepstorff, “A class of anomaly-free gauge theories,” arXiv:hep-th/0005079; “Towards a unified theory of gauge and Yukawa interactions,” arXiv:hep-ph/0006065; “Extra dimensions: Will their spinors play a role in the standard model?,” arXiv:hep-th/0310092; F. D. Smith, “From sets to quarks: Deriving the standard model plus gravitation from simple operations on finite sets,” arXiv:hep-ph/9708379. J.S.R. Chisholm and R.S. Farwell, *J. Phys. A: Math. Gen.* **20**, 6561 (1987); **33**, 2805 (1999); **22**, 1059 (1989); J.S.R. Chisholm, *J. Phys. A: Math. Gen.* **35**, 7359 (2002); *Nuov. Cim.* **A82**, 145 (1984); 185; 210; “Properties of Clifford Algebras for Fundamental Particles”, in *Clifford (Geometric) Algebras*, W. Baylis ed., Birkhauser (1996), Chapter 27, pp. 365–388. J.P. Crawford, *J. Math. Phys.* **35**, 2701 (1994); in *Clifford (Geometric) Algebras*, W. Baylis ed., Birkhauser (1996), Chapters 21–26, pp. 297–364; *Class. Quant. Grav.* **20**, 2945 (2003); A. Garrett Lisi, arXiv:gr-qc/0511120.